

Stanford Artificial Intelligence Laboratory
Memo AIM-222

AUGUST 1974

Computer Science Department
Report No. STAN-CS-74-46 7

Checking Proofs in the Metamathematics of First Order Logic

by

Mario Aiello
Richard W. Weyhrauch

Research sponsored by

Advanced Research Projects Agency
ARPA Order No. 2494

COMPUTER SCIENCE DEPARTMENT
Stanford University



STAN-CS-74-467

CHECKING PROOFS IN THE METAMATHEMATICS OF FIRST ORDER LOGIC

by
Mario Aiello
and
Richard W. Weyhrauch

Abstract:

This is a report on some of the first experiments of any size carried out using the new first order proof checker FOL. We present two different first order axiomatizations of the metamathematics of the logic which FOL itself checks and show several proofs using each one. The difference between the axiomatizations is that one defines the metamathematics in a many sorted logic, the other does not.

Authors' addresses

M. Aiello, Istituto di Scienze dell'Informazione, Universita' di Pisa, corso Italia 40, 56100 Pisa, Italy;

R. Weyhrauch, A.I. Lab., Computer Science Dept., Stanford University, Stanford, California 94305. or
Weyhrauch @SU-AI

This research is supported (in part) by the Advanced Research Project Agency of the Office of the Secretary of Defense (DAHC 15-73-G-0435).

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Advanced Research Project Agency, the U.S. Government.

Reproduced in USA. Available from the National Technical Information Service, Springfield, Virginia 22151.

Checking metamathematical proofs

TABLE OF CONTENTS

1 INTRODUCTION	1
2 THE AXIOM SYSTEM	2
2.1 The sorts	2
2.2 The domain of representation of the metamathematics	3
2.2.1 Formulas and terms	4
2.2.2 Rules of inference, deductions and the notion of provability	6
2.3 The main proof in the many sorted logic	7
2.4 Another axiomatization	7
3 THE PROOFS	8
3.1 A look at sorts	8
3.2 The unify and tautology commands	8
4 CONCLUSION	11
Appendix 1 THE AXIOMS IN THE MANY SORTED LOGIC	12
1.1 Natural numbers	12
1.2 The set of symbols	12
1.3 Strings	12
1.4 Formulas	13
1.5 Sequences	13
1.6 Free and bound variables and the substitution	14
1.7 Rules of inference	15
1.8 Deduction	16
Appendix 2 THE AXIOMS IN THE LOGIC	18
2.1 Natural numbers	18
2.2 The set of symbols	18

1000

1000

1000

Checking metamathematical proofs'	1
2.3 Strings	18
2.4 Sequences	19
2.5 Formulas	20
2.6 Free and bound variables and the substitution	20
2.7 Rules of inference	22
2.8 Deduction	23
Appendix 3 THE PROOF OF "IF f IS A WFF ALSO $\forall x.f$ IS A WFF"	25
3.1 FOL commands and printout in the many sorted logic	25
3.2 FOL commands in the earlier axiomatization	25
3.3 Printout of the proof in the earlier axiomatization	26
Appendix 4 THE PROOF OF THE EQUIVALENCE BETWEEN SBV AND SBT FOR VARIABLES	28
4.1 FOL commands in the many sorted logic	28
4.2 Printout of the proof in the many sorted logic	29
4.3 FOL commands in the earlier axiomatization	30
4.4 Printout of the proof in the earlier axiomatization	31
Appendix 5 THE PROOF THAT UNIVERSAL QUANTIFIER CAN BE INTERCHANGED	35
5.1 FOL commands for the main lemma in the many sorted logic	35
5.2 Printout of the proof in the many sorted logic	35
5.3 FOL commands for the theorem in the many sorted logic	36
5.4 Printout of the proof of the theorem in the many sorted logic	38
5.5 FOL commands for the main lemma in the earlier axiomatization	41
5.6 Printout of the proof of the main lemma in the second axiomatization	42
5.7 FOL commands in the earlier axiomatization	43
5.6 Printout of the proof in the earlier axiomatization	45

Checking metamathematical proofs

SECTION 1 INTRODUCTION

This paper represents a first attempt at the axiomatization of the metamathematics of a first order theory and at using the new proof checker FOL (First Order Logic). The logic which FOL checks is described in detail in the user manual for this program, Weyhrauch and Thomas 1974. It is based on a system of natural deduction described in Prawitz 1965, 1970.

Our motivation in axiomatizing the metamathematics of FOL was the desire to work on an example which could be used as a case study for projected features of FOL and, at the same time, had independent interest with respect to representing the proofs of significant mathematical results to a computer.

The eventual ability to clearly express the theorems of mathematics to a computer will require the facility to state and prove theorems of metamathematics. There are several clear examples:

a. *Axiom schemas*. How exactly do we express that

$$P(0) \wedge \forall n.(P(n) \supset P(n+1)) \supset \forall n.P(n)$$

is an axiom schema? We need to say: "If for any first order sentence P with one free variable y we denote by P(n) the formula obtained from P by substituting n for y assuming n is free for y in P, then the sentence

$$P(0) \wedge \forall n.(P(n) \supset P(n+1)) \supset \forall n.P(n)$$

is an axiom of arithmetic".

b. *Theorem schemas*. The following kind of "theorem" is sometimes seen in set theory books

$$\forall x_1 \dots x_n \in S. \exists T. \forall u. (\langle x_1, \dots, x_n \rangle \in T \supset \exists y. (\langle x_1, \dots, x_n, y \rangle \in S)).$$

It asserts the existence of some particular projection of n+1-tuples. In its usual formulation this is not a theorem of set theory at all, but a metatheorem which states that, for each n, the above sentence is a theorem. We do not know of any implementation of first order logic capable of expressing the above notion in a straightforward way.

c. *Subsidiary deduction rules*. Below we show how to prove that if there is a proof of $\forall x y. WFF$ then there is also a proof of $\forall y x. WFF$, where WFF is any well formed formula. We chose this task because it seemed simple enough to do, and is a theorem which may actually be used. The use of metatheorems as rules of inference by means of a reflection principle will be discussed in a future memo by Richard Weyhrauch. Eventually we hope to check some more substantial metamathematical theorems.

d. *Interesting mathematical theorems*. We present two examples. The first is any theorem about finite groups. The notion of finite group cannot be defined in the usual first order language of group theory. Thus many "theorems" are actually metatheorems, unless you axiomatize groups in set theory. The second theorem is the "duality principle" in projective geometry.

Checking metamathematical proofs

$sy\ sy1\ sy2\ sy3\ sy4\ sy5\ sy6 \in SYM$ (SYM are logical symbols)
 $np\ np1\ np2\ np3\ np4\ np5\ np6 \in N_PLCSYM$ (N_PLCSYMs are symbols which have an *arity*)
 $fn\ fn1\ fn2\ fn3\ fn4\ fn5\ fn6 \in OPCONST$, (OPCONSTs are function symbols)
 $P\ P1\ P2\ P3\ P4\ P5\ P6 \in PRDCONST$; PREDCONSTs are predicate symbols)

the partial order between these sorts is defined by the following FOL declarations:

```

MG SEQ      ≥ { STRING , PROOFTREE };
MC PROOFTREE ≥ { FORM };
MG STRING   ≥ { TERM , FORM , ATOM , VARSTRING };
MG TERM     ≥ { INDVAR };
MG FORM     ≥ { ELF , SENTCONST , PREDPAR0 , AXIOM , BEW };
MG BEW      ≥ { AXIOM };
MG ATOM     ≥ { INDCONST , SENTCONST , SYM , INTEGER , N_PLCSYM ,
               INDPAR , INDVAR , AUXSIGN , PREDCONSTB , PREDPAR0 };
MG INDCONST ≥ { NUMERAL };
MG SYM      ≥ { QUANT , SENTCONN };
MG N_PLCSYM ≥ { PREDCONST , OPCONST , PREDPAR };
  
```

Sorts are always predicates with one argument. The declaration

$M\ C\ SORT1 \geq \{ SORT2 , SORTn \}$

should be read as SORT1 is more general than SORT2,...,SORTn and corresponds to the implicit axioms

$\forall g. SORT1(g) \Rightarrow SORT1(g),$

$\forall g. SORTn(g) \Rightarrow SORT1(g)$

The first declaration, for instance, says that strings and derivations are particular sequences of formulas. Strings are in fact sequences of length 1 and derivations are those sequences satisfying the predicate PROOFTREE.

Section 2.2 The domain of representation of the metamathematics

The basic notions of the metamathematics of first order logic have been axiomatized in terms of strings and sequences of strings. The primitive functions on them are concatenation (*c* for strings, *cc* for sequences) and selectors (*car*, *cdr* for strings and *scar*, *scdr* for sequences) *c* and *cc* are infix operators.

SECTION 2 THE AXIOM SYSTEM

In this section we present two **axiomatizations** of the metamathematics of first order logic, The main difference between them is that one is done in a many sorted **first** order logic and the other not. These **axiomatizations** represent an attempt at experimenting with proofs about properties of formulas and deductions. No effort has been spent on guaranteeing that the axioms are independent. It would not only have been uninteresting but also contrary to our basic philosophy. We wish to find axioms which naturally reflect the relevant notions. At the moment this **axiomatization** is far from being in its final form. Neither the extent of the notions involved nor the best way of expressing them is considered settled.

Section 2.1 The sorts

The sorts we have defined correspond to the basic notions of the metamathematics i.e. terms, formulas, individual variables, **logical** symbols, function symbols etc. and to the notions of the domains (strings and -sequences of strings) in which the axiomatization has been defined. FOL (see Weyhrauch and Thomas 1974) allows the declaration of variables to be of a certain sort. In the formulas appearing in this paper the following declarations are assumed:

g g1 g2 g3 g4 g5 g6	range over the most general sort
sq sq1 sq2 sq3 sq4 sq5 sq6 $\in$ SEQ	(SEQs are sequences of strings)
pf pf1 pf2 pf3 pf4 pf5 pf6 $\in$ PROOFTREE	(PROOFTREEs are sequences representing derivations in FOL)
s s1 s2 s3 s4 s5 s6 $\in$ STRING	(STRINGs are strings)
t t1 t2 t3 t4 t5 t6 $\in$ TERM	(TERMs are strings representing terms)
x x1 x2 x3 x4 x5 x6 $\in$ INDVAR	(INDVARs are strings representing individual variables)
e1 e11 e12 e13 e14 e15 e16 $\in$ E L F	(ELFs are strings representing elementary formulas)
f f1 f2 f3 f4 f5 f6 $\in$ FORM	(FORMs are well formed formulas)
th th1 th2 th3 th4 th5 th6 $\in$ BEW	(BEWs are theorems of a first order theory)
A A1 A2 A3 A4 A5 A6 $\in$ AXIOM	(AXIOMs are axioms of a particular theory)
c0 c1 c2 c3 c4 c5 c6 $\in$ INDCONST	(INDCONSTs are individual constants)
a a1 a2 a3 a4 a5 a6 $\in$ ATOM	(ATOMs are the individual constituents of a string)
n n1 n2 n3 n4 n5 k $\in$ INTEGER	(INTEGERs are integers)
nc nc1 nc2 nc3 nc4 nc5 nc6 $\in$ NUMERAL	(NUMERALs are numerals)

The properties of wffs relevant to our theory have been defined by the predicates FR, FRN, GEB and SBT. FR(x,f) is true iff the variable x has at least one free occurrence in the wff f, while FRN(x,n,f) and GEB(x,n,f) are respectively true when the variable x occurs free or bound at the place n in the formula f. These predicates are defined in appendix 1.6. In addition, some generalized selector functions are defined, which evaluate the first or the k-th free occurrence of a variable in a wff, or the number of its free occurrences. The predicate SBT is then defined. It axiomatizes the notion of substitution of a term for any free occurrence of a variable in a wff.

$$\begin{aligned} \forall x \, t \, f1 \, f2. (SBT(x,t,f1,f2) = & \\ \forall n1 \, n2. ((n2 = (\text{numfreeocc}(x,n1,f1) * (\text{len}(t)-1)) + n1) \supset & \\ ((\neg \text{INDVAR}(n1 \, \text{gl} \, f1) \supset (n1 \, \text{gl} \, f1) = (n2 \, \text{gl} \, f2)) \wedge & \\ (\text{INDVAR}(n1 \, \text{gl} \, f1) \supset ((\text{FRN}(x,n1,f1) \supset \text{SUBT}(t,f2,n2)) \wedge & \\ (\neg \text{FRN}(x,n1,f1) \supset \text{INVART}(n1,f1,n2,f2)))))) \wedge & \\ \forall t \, f2 \, n2. (\text{SUBT}(t,f2,n2) = \forall x2 \, k. ((k \, \text{gl} \, t) = x2 \supset \text{FRN}(x2,n2-(\text{len}(t)-k),f2))))), & \\ \forall n \, f1 \, n1 \, f2. (\text{INVART}(n,f1,n1,f2) = ((\text{GEB}(n1 \, \text{gl} \, f2,n1,f2) = \text{GEB}(n \, \text{gl} \, f1,n,f1)) \wedge & \\ (\text{FRN}(n1 \, \text{gl} \, f2,n1,f2) = \text{FRN}(n \, \text{gl} \, f1,n,f1)) \wedge (n1 \, \text{gl} \, f2) = (n \, \text{gl} \, f1))) & \end{aligned}$$

In the previous definition, n1 is any position in the string f1 and n2 is the corresponding position in f2. The auxiliary predicate SUBT states that the variables appearing in the term t substituted for a free occurrence of the variable x are still free. INVART defines which properties of f1 are still true for f2. If the term t is a variable, then SBT reduces to SBV:

$$\begin{aligned} \forall x1 \, x2 \, f1 \, f2. (\text{SBV}(x1,x2,f1,f2) = & \\ \forall n1. ((\neg \text{INDVAR}(n1 \, \text{gl} \, f1) \supset (n1 \, \text{gl} \, f1) = (n1 \, \text{gl} \, f2)) \wedge & \\ (\text{INDVAR}(n1 \, \text{gl} \, f1) \supset ((\text{FRN}(x1,n1,f1) \supset \text{FRN}(x2,n1,f2)) \wedge & \\ (\neg \text{FRN}(x1,n1,f1) \supset \text{INVARV}(n1,f1,f2)))))) \wedge & \\ \forall n \, f1 \, f2. (\text{INVARV}(n,f1,f2) = ((\text{GEB}(n \, \text{gl} \, f2,n,f2) = \text{GEB}(n \, \text{gl} \, f1,n,f1)) \wedge & \\ (\text{FRN}(n \, \text{gl} \, f2,n,f2) = \text{FRN}(n \, \text{gl} \, f1,n,f1)) \wedge (n \, \text{gl} \, f2) = (n \, \text{gl} \, f1))), & \end{aligned}$$

The proof of the equivalence of SBT and SBV when t is a variable is very simple. It is based on the fact that n2 coincides with n1 when the term t has length 1 (see appendix 4). The function sbt (sbv) evaluates to the string representing the result of substituting a term (variable) for every free occurrence of a variable in a given wff. sbt and sbv are defined from the predicates SBT and SBV as follows:

$$\begin{aligned} \forall x \, t \, f1 \, f2. (\text{SBT}(x,t,f1,f2) = \text{sbt}(x,t,f1,f2)) \\ \forall x1 \, x2 \, f1 \, f2. (\text{SBV}(x1,x2,f1,f2) = \text{sbv}(x1,x2,f1,f2)) \end{aligned}$$

The problem of finding the best way of defining functions in FOL is crucial in the axiom system given in this paper: a uniform way has not been followed. In defining the substitution we are interested in properties of the functions sbt and sbv and in drawing conclusions from the fact that a substitution has been made. It is thus useful to have a predicate which defines the relation between formulas before and after a substitution instead of inferring it from the definitions of the functions (stated for example as a system of equations, as in Kleene 1952). One of the motivations of the present experiment was to explore different ways of defining functions. We do not yet have enough examples of proofs to make a clear statement about this matter.

2.2.1 Formulas and terms

Formulas and terms are represented by the string of symbols appearing in **them**. Terms are defined **recursively** as strings which either represent an individual variable or can be decomposed into **$n+1$ substrings representing** a function symbol of **arity n** , followed by **n terms**. The two predicates defining terms are:

TERMSEQ(0,LAMBDA)

$\forall s. (\text{TERM}(s) \equiv \text{INDVAR}(s) \vee \exists n \text{ fn}. (\text{fn} = \text{car}(s) \wedge n = \text{arity}(\text{fn}) \wedge \text{TERMSEQ}(n, \text{cdr}(s))))$

$\vee n \ s. (\text{TERMSEQ}(n, s) \equiv ((\text{car}(s) = \text{LPARSYM}) \wedge ((\text{len}(s) \text{ gl } s) = \text{RPARSYM}) \wedge \exists n1. (\text{TERM}(\text{substring}(s, 2, n1)) \wedge \text{TERMSEQ}(n-1, \text{substring}(s, n1+1, \text{len}(s)-1))))))$

where the function **substring(s, m, n)** (see appendix 1.3) returns the substring of s starting from its m -th element and ending with the n -th. **len(s)** computes the length of s and **(n gl s)** selects the n -th element of s .

Well formed formulas (wffs) are represented as strings which either are elementary formulas (**defined** by the predicate **ELF**) or can be partitioned into substrings for formulas and logical connectives. Formulas are defined by:

$\forall s. (\text{ELF}(s) \equiv (s = \text{FALSESYM} \vee \text{PREDPARO}(s) \vee \exists n \ P. (P = \text{car}(s) \wedge n = \text{arity}(P) \wedge \text{TERMSEQ}(n, \text{cdr}(s)))))$,

$\forall s. (\text{FORM}(s) \equiv (\text{ELF}(s) \vee \exists x \ f. (s = (x \text{ gen } f) \vee s = (x \text{ ex } f)) \vee \exists f1 \ f2. (s = (f1 \text{ dis } f2) \vee s = (f1 \text{ con } f2) \vee s = (f1 \text{ impl } f2)) \vee \exists f. s = \text{neg}(f)))$;;

gen is the infix operator that maps its arguments x and f into the string **(FORALLSYM $c \ x$) $c \ f$ representing** the well formed formula $\forall x.f$. The operator **ex** is used for the existential quantifier. **dis**, **con** and **impl** are the infix operators for the disjunction, conjunction and implication of two formulas. Finally, **neg** is the operator which maps a formula into its negation.

We could possibly represent wffs as structured objects (lists, trees, etc.) which contain all the **information** about the structure of the formula and do not require any parsing. This approach **amounts to** axiomatizing metamathematics in terms of the abstract syntax of first order logic, instead of strings of symbols. Both of these possibilities should be explored. We have chosen the first alternative because:

1) It is the most traditional, i.e. **metamathematics**, as it appears in logic books, is usually stated in terms of strings.

2) **Axioms in** terms of abstract syntax are simply theorems of the theory expressed in terms of strings. Thus the **two** representations look substantially the same with respect to "high level" theorems.

3) Ill-formed formulas can be mentioned. This is of course impossible in an axiomatization in terms of the abstract syntax.

The **main** theorem we have proved in this axiomatization of the metamathematics states **that if $\forall x y.wff$ is provable in some theory, then $\forall y x.wff$ is also provable.** We have chosen this theorem because, even if very simple, it involves basic notions of provability, substitution and universal quantification. Its proof is found in appendices 5.1-2. The theorem depends on the first three lines of the proof. The first step is a lemma stating that $\forall x wff.sbt(x,x,f)=wff$, i.e. substituting a variable x for any free occurrence of x in wff doesn't change that wff . Steps two and three give simple facts about sequences. The theorem is then proved by instantiating two other lemmas: 1) if $\forall x.wff$ is a **theorem, then wff is also a theorem**; 2) if wff is provable, then x cannot be free in the dependencies of the proof of wff and so $\forall x.wff$ is provable. This is of course true only for theories with no free variables in their axioms.

The only property of the inference rules used in this proof involves universal quantification. The restriction on the applicability of the $\forall$ -introduction rule is, that the variable to be **universally** quantified in a wff must not appear free in any of its dependencies. This restriction is reflected in our axiomatization by the predicate APGENI. In this proof APGENI is satisfied because if wff is provable, its dependencies are axioms with no free variables.

The following is an informal proof of the above theorems. If $\forall x.wff$ is provable, then there **is a proof tree pf** whose first string is $\forall x.wff$. The sequence $(\forall x.wff) \text{ cc } pf$ is still a proof tree. It is obtained by applying the $\forall$ -elimination rule. The application of this rule doesn't add any dependency to the proof tree. As its only dependencies are axioms, it follows from the definition of BEW that wff **is a theorem**. On the other hand, if wff is a theorem there exists a **proof tree pf** whose first element is wff . By applying the $\forall$ -introduction rule to pf we obtain the proof tree $(\forall x.wff) \text{ cc } pf$. This rule is applicable since theorems have no free variables in their dependencies. It follows that $\forall x.wff$ is a theorem. If $\forall x y.wff$ is provable then $\forall x.wff$ and wff are provable using the first lemma. Finally, we can quantify first over x and then over y , obtaining $\forall y x.wff$ as a theorem.

Section 2.4 Another axiomatization

A different axiomatization has been given in an earlier version of FOL where there was no facility for creating sorts. We present it here as we want to do some comparisons between proofs, and discuss some of the features of FOL. Some differences between the two axiomatizations are due to the new features available in FOL. They will be discussed in the next section. Here we only **discuss** the difference between the definition of formulas and terms. The list of all the axioms can be found in appendices 2 f 8.

In this axiomatization, formulas and terms are still represented as the string of the symbols appearing in them. They are defined as strings that can be decomposed into a sequence of substrings recording the construction of that formula or term from elementary formulas and individual variables, according to the usual formation rules (see appendix 2.5 for the list of axioms). These sequences are defined by the predicate TERMSEQ for terms and FRR for wffs. A sequence satisfies the predicate TERMSEQ if it represents the history of the construction of its first element (the term to be defined), starting from symbols, functions and individual variables. Similarly, a string is a wff if there exists a sequence which satisfies the predicate FRR and represents the history of the construction of that wff from elementary formulas and the logical connectives.

Checking **metamathematical** proofs

2.2.2 Rules of inference, deductions and the notion of provability

The rules of inference are defined by the predicates in appendix 1.7. The rules with one premise, are expressed by means of a binary predicate whose arguments are two sequences of wffs (sq, pf) which satisfy PROOFTREE. The predicate is true iff pf is the **schr** of sq and the first element of sq is a wff obtained by applying that particular deduction rule to the first wff of pf. The rules with more antecedents are defined in a similar way.

Derivations are recursively defined as sequences of wffs which either are a single wff or are obtained from one or more derivations by applying one of the deduction rules. The recursion is implicitly stated by saying that there exist objects of sort PROOFTREE which satisfy one of the predicates defining the rules of inference. These sequences represent **the linearization** of a **deduction-tree** and are defined as follows:

$$\begin{aligned} \forall sq. (PROOFTREE(sq) \equiv & \\ & (FORM(sq) \vee \\ & \exists pf. (ORI(sq, pf) \vee ANDE(sq, pf) \vee FALSEE(sq, pf) \vee NOTI(sq, pf) \vee NOTE(sq, pf) \vee IMPLI(sq, pf)) \vee \\ & \exists pf \ x \ t. (GENI(sq, pf, x, t) \vee GENE(sq, pf, x, t) \vee EXI(sq, pf, x, t)) \vee \\ & \exists pf \ 1 \ pf2. (ANDI(sq, pf \ 1, pf2) \vee FALSEI(sq, pf \ 1, pf2) \vee IMPE(sq, pf \ 1, pf2)) \vee \\ & \exists pf \ 1 \ pf2 \ x \ 1 \ x2. EXE(sq, pf \ 1, pf2, x \ 1, x2) \vee \\ & \exists pf \ 1 \ pf2 \ pf3. ORE(sq, pf \ 1, pf2, pf3)) \vee \vee \end{aligned}$$

A sequence of wffs is a **prooftree** if either it consists of a single wff or one of the following alternatives holds: there exists another prooftree and a one premise deduction rule has been applied; there exist two prooftrees and one of the two premises rules has been applied; finally, there are three prooftrees and the predicate defining the $\vee$ -elimination rule is true. Note that the root of a prooftree is not necessarily a **theorem** in a given **theory**. A predicate **DEPEND** has been defined which is true if a given wff is a **dependence** for the root of a prooftree. The axioms about **DEPEND** allows to decide all the dependencies of a prooftree.

Since some of the deduction rules (the implication introduction, for instance) eliminate dependencies, not all the leaves of a prooftree pf are dependencies for a wff f such that **f=schr(pf)**. The predicate **DEPEND** is true only for those **leaves** of the prooftree which the formula f actually depends on, its definition is shown in appendix 1.8. The axioms **DEPEND** state which dependencies do not change by applying the deduction rules and are transferred from one prooftree to the other. The axioms **NDEPEND** state which rules discharge dependencies in a given prooftree.

Using this notion of dependence, the provability of a formula in a theory is defined as follows:

$$\forall f. (BEW(f) \equiv \exists sq. (PROOFTREE(sq) \wedge f = schr(sq) \wedge \forall f \ 1. (DEPEND(sq, f \ 1) \supset AXIOM(f \ 1))));$$

A wff f is a **theorem** in a given theory if there exists a prooftree whose first element is f and whose only dependencies are axioms in that theory. We have limited our attention to theories in which axioms have no free variables. This property is defined by the axiom:

$$\forall x \ f. (AXIOM(f) \supset \neg FR(x, f));;$$

Section 2.3 The **main** proof in **the many** sorted logic

statements. Hence, the statements produced by them have quantifiers as main symbols or it is necessary to introduce a quantifier to proceed in the proof. After the right introductions or eliminations have been done to them, the tautology commands are used again. This process is iterated until the completion of the proof.

The command UNIFY decides if a given wff can be obtained by instantiation of quantified variables or introduction of them for free occurrences of variables or terms in a second wff. The code for this command has been written by Ashok Chandra and is still in an experimental stage. In the proofs presented here, this command has been essentially used for the simultaneous introduction of the existential quantifier. As an example, consider the following assumption:

```
1  $\forall x.(P(x) \Rightarrow (Q(f) \wedge \forall t.R(t)))$  (1) ASSUME
```

the command

```
unify  $\exists x.(P(x) \Rightarrow \exists f.(Q(f) \wedge R(g(t))))$ ,1;
```

deduces in a single step

```
2  $\exists x.(P(x) \Rightarrow \exists f.(Q(f) \wedge R(g(t))))$  (6) UNIFY 1
```

A good example of a combined use of these features is found in appendix 3.3:

```
19 FRR((x1 gen f) cc SQ) (SEQUENCE((x1 gen f) cc SQ) ^ (((x1 gen f) cc U) ^
  SLAMBDA ^ (ELF(scar((x1 gen f) cc SQ)) v (FRR(scdr((x1 gen f) cc SQ)) ^
   $\exists s1 s2. (STRING(s1) \wedge (STRING(s2) \wedge ((scar((s1 gen f) cc SQ) = NEG(s1) \wedge$ 
  find(1,s1,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 dis s2) ^
  find(2,s1 c s2,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 con s2) ^
  find(2,s1 c s2,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 impl s2) ^
  find(2,s1 c s2,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 gen s2) ^ (INDVAR(s1) ^
  find(1,s2,scdr((x1 gen f) cc SQ))) v (scar((x1 gen f) cc SQ) = (s1 ex s2) ^ (INDVAR(s1) ^
  find(1,s2,scdr((x1 gen f) cc SQ))))))))) ---  $\forall E$  WFF1(x1 gen f) cc SQ

20 STRING(x1) ^ (STRING(f) ^ ((scar((x1 gen f) cc SQ) = NEG(x1) ^
  find(1,x1,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (x1 dis f) ^
  find(2,x1 c f,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (x1 con f) ^
  find(2,x1 c f,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (x1 impl f) ^
  find(2,x1 c f,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (x1 gen f) ^ (INDVAR(x1) ^
  find(1,f,scdr((x1 gen f) cc SQ))) v (scar((x1 gen f) cc SQ) = (x1 ex f) ^ (INDVAR(x1) ^
  find(1,f,scdr((x1 gen f) cc SQ))))))))) (1 2 3 4 5 6 7 8 11) --- TAUTEQ 1:19

21  $\exists s1 s2. (STRING(s1) \wedge (STRING(s2) \wedge ((scar((s1 gen f) cc SQ) = NEG(s1) \wedge$ 
  find(1,s1,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 dis s2) ^
  find(2,s1 c s2,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 con s2) ^
  find(2,s1 c s2,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 impl s2) ^
  find(2,s1 c s2,scdr((x1 gen f) cc SQ))) v ((scar((x1 gen f) cc SQ) = (s1 gen s2) ^ (INDVAR(s1) ^
  find(1,s2,scdr((x1 gen f) cc SQ))) v (scar((x1 gen f) cc SQ) = (s1 ex s2) ^ (INDVAR(s1) ^
  find(1,s2,scdr((x1 gen f) cc SQ))))))))) (1 2 3 4 5 6 7 8 11) --- UNIFY 20
```

Line 19 is the instantiation of an axiom. Line 20 is generated by the command,

```
TAUTEQ 19: #2#2#2#2#2#1#1[s1←f : s2←x1]1:19;
```

note how the use of the **FOL** subpart designators allows us to mention the desired subpart of 19, without **having** to retype it. In addition we can do the appropriate substitutions. **Line 21** is just a use of UNIFY:

UNIFY 19:02*2*2*2*2 20;

Because we can mention the conclusion, without writing it down explicitly, the amount of typing necessary is severely reduced. Without UNIFY, line 21 would have required two **3-introductions** and the commands would have been:

31 20 xl+-sl OCC 1,2,3,4,7,8,11,12,15,16,19,20,23,24;

31 20 f ←s2 occ 1,5,9,13,17,18,21,22;

We do not enter into a detailed discussion of the command UNIFY. It is our **intension** to do it elsewhere. It should be thought of as the routine which handles quantifiers in "simple" inferences. As seen above, the saving to a user can be large.

SECTION 4 CONCLUSION

The desire to represent mathematics in a computer in a feasible way certainly requires the facility to discuss metamathematical notions. The axiomatization presented here only treats the syntactic part of the problem. Any mention of the models involved needs the addition of set theory to the axiomatization. However, it is clear from the simple theorems we proved that any practical system needs more extensive features even to do a satisfactory job of writing down the theorems we might want.

An important point for future work is how (in a practical way) to use these theorems. Consider for instance:

$$\forall x_1 x_2 f. (\text{BEW}(x_1 \text{ gen } (x_2 \text{ gen } f)) \supset \text{BEW}(x_2 \text{ gen } (x_1 \text{ gen } f)))$$

What we mean by reflection principle is a rule of FOL which says:

$$\frac{\text{//BEW}(f) \quad \text{//in meta FOL}}{\text{//}f} \quad \text{// in FOL}$$

That is, if in the axiomatization of the metamathematics of FOL, we can prove the existence of an FOL proof of f , then we can assert f in FOL. Suppose we have a proof in FOL of $\forall x y. wff$. Then instantiating the above theorem gives us

$$\text{BEW}(x \text{ gen } (y \text{ gen } wff)) \supset \text{BEW}(y \text{ gen } (x \text{ gen } wff))$$

Since we started with a proof of $\forall x y. wff$ in FOL and BEW represents the proof predicate for FOL, we can conclude $\text{BEW}(x \text{ gen } (y \text{ gen } wff))$. Using modus ponens we get $\text{BEW}(y \text{ gen } (x \text{ gen } wff))$, and using the above rule we can conclude $\forall y x. wff$ in FOL.

The exact form of such a rule requires more examples of proofs and is one of the main reasons for doing the example in the memo. It is not just a proof checking exercise, but a case study for fundamental questions of representing mathematical information in a computer. Using metamathematics also prepares the way for more comprehensive systems which can formally discuss how they reason. That is exactly what the metamathematics is good for.

APPENDIX I

THE AXIOMS IN THE MANY SORTED LOGIC

1.1 Natural numbers

AXIOM NUMB:

$$\begin{aligned}
& \forall n1 \ n2 \ n3. (n1 = n2 \supset (n1 = n3 \supset n2 = n3)), \\
& \forall n1 \ n2. (n1 = n2 \supset succ(n1) = succ(n2)), \\
& \forall n1. 0 \neq succ(n1), \\
& \forall n1 \ n2. (succ(n1) = succ(n2) \supset n1 = n2), \\
& \forall n1. n1 + 0 = n1, \\
& \forall n1 \ n2. n1 + succ(n2) = succ(n1 + n2), \\
& \forall n1. n1 * 0 = 0, \\
& \forall n1 \ n2. n1 * succ(n2) = (n1 * n2) + n1 ;;
\end{aligned}$$

A X I O M INDCT:

$$(F(0) \wedge \forall n. (F(n) \supset F(n+1))) \supset \forall n. F(n) ;;$$

AXIOM DEFN:

$$\begin{aligned}
& \forall n. (succ(n)-1) = n, \\
& \forall n1 \ n2. succ(n1) - n2 = n1 - (n2 - 1), \\
& \forall n1 \ n2 \ n3. (n1 < n2 \equiv \exists n3. (n3 \neq 0 \wedge n1 + n3 = n2)), \\
& \forall n1 \ n2. (n1 \leq n2 \equiv (n1 < n2) \vee (n1 = n2)), \\
& \forall n1 \ n2. (n2 > n1 \equiv n1 < n2), \\
& \forall n1 \ n2. (n2 \geq n1 \equiv n1 \leq n2) ;;
\end{aligned}$$

1.2 The set of symbols

AXIOM SYM:

$$\begin{aligned}
& \forall s. (SYM(a) \equiv a = LPARSYM \vee a = RPARSYM \vee a = ORSYM \vee a = ANDSYM \vee a = IMPSYM \vee \\
& \quad a = FALSESYM \vee a = FORALLSYM \vee a = EXISTSYM) ;;
\end{aligned}$$

1.3 Strings

AXIOM STRING:

$$\begin{aligned}
& \forall s. s = car(s) \circ cdr(s), \\
& \forall s1 \ s2. (s1 = LAMBDA \supset car(s1 \circ s2) = car(s2)), \\
& \forall s1 \ s2. (s1 \neq LAMBDA \supset car(s1 \circ s2) = car(s1)), \\
& \forall s1 \ s2. (s1 = LAMBDA \supset cdr(s1 \circ s2) = cdr(s2)), \\
& \forall s1 \ s2. (s1 \neq LAMBDA \supset cdr(s1 \circ s2) = cdr(s1)), \\
& \forall s. (s \circ LAMBDA = LAMBDA \circ s), \\
& \forall s. s \circ LAMBDA = s, \\
& \forall s1 \ s2 \ s3. (s1 \circ (s2 \circ s3) = (s1 \circ s2) \circ s3), \\
& \forall a. (len(a) = 1 \vee a = LAMBDA), \\
& \forall s. len(s) \geq 0, \\
& \forall s1 \ s2. len(s1 \circ s2) = len(s1) + len(s2), \\
& \forall s. (len(s) = 1 \supset ATOM(s)), \\
& \forall s. \exists gl \ s = LAMBDA,
\end{aligned}$$

VS. $1 \text{ gls} = \text{car}(s)$,
 Vs n. $((n > 1) \supset ((n \text{ gl } s) = ((n-1) \text{ gl } \text{cdr}(s)))) ; ;$

AXIOM SUBSTRDEF:

$\forall n1 \ n2 \ s1 \ s2. \quad (\text{SUBSTP}(s1, s2, n1, n2) \equiv (\text{len}(s2) = n2 - n1 + 1 \wedge (\forall n. (n \geq n1 \wedge n \leq n2 \supset n \text{ gl } s1 = (n - n1 + 1) \text{ gl } s2))))$,
 $\forall n1 \ n2 \ s1 \ s2. \quad (\text{SUBSTP}(s1, s2, n1, n2) \equiv \text{substring}(s1, n1, n2) = s2)$,
 $\forall s1 \ s2. \quad (\text{SUBS}(s1, s2) \equiv \exists n1 \ n2. \text{SUBSTP}(s1, s2, n1, n2)) ; ;$

The value of $\text{substring}(s1, n1, n2)$ is the substring of $s1$ whose first element is the $n1$ th element of $s1$ and whose last element is the $n2$ th element.

AXIOM DISEQ:

$\forall g1 \ g2. (\neg(g1 = g2) \equiv g1 \neq g2) ; ;$

AXIOM EQS:

$\forall s1 \ s2. (\forall n. (n \text{ gl } s1 = n \text{ gl } s2) \equiv s1 = s2) ; ;$

AXIOM COMP:

$\forall f. \quad e(f) = (\text{LPARSYM } c \ f) \ c \ \text{RPARSYM}$,
 $\forall f1 \ f2. \quad f1 \ \text{dis} \ f2 = (e(f1) \ c \ \text{ORSYM}) \ c \ e(f2)$,
 $\forall f1 \ f2. \quad f1 \ \text{impl} \ f2 = (e(f1) \ c \ \text{IMPSYM}) \ c \ e(f2)$,
 $\forall f. \quad \text{neg}(f) = (f \ \text{impl} \ \text{FALSESYM})$,
 $\forall f1 \ f2. \quad f1 \ \text{con} \ f2 = (e(f1) \ c \ \text{ANDSYM}) \ c \ e(f2)$,
 $\forall x \ f2. \quad x \ \text{gen} \ f2 = (\text{FORALLSYM } c \ x) \ c \ f2$,
 $\forall x \ f2. \quad x \ \text{ex} \ f2 = (\text{EXISTSYM } c \ x) \ c \ f2 ; ;$

1.4 Formulas

AXIOM TERM:

$\forall n \ s. \quad (\text{TERMSEQ}(n, s) \equiv (\exists n1. (\text{TERM}(\text{substring}(s, 1, n1)) \wedge \text{TERMSEQ}(n-1, \text{substring}(s, n1+1, \text{len}(s)))))$,
 vs. $(\text{TERM}(s) \equiv \text{INDVAR}(s) \vee \exists n \ \text{fn}. (\text{fn} = \text{car}(s) \wedge n = \text{arity}(\text{fn}) \wedge \text{TERMSEQ}(n, \text{cdr}(s)))) ; ;$

AXIOM WFF:

vs. $(\text{ELF}(s) \equiv (s = \text{FALSESYM} \vee \text{PREDPAR0}(s) \vee \exists n \ P. (P = \text{car}(s) \wedge n = \text{arity}(P) \wedge \text{TERMSEQ}(n, \text{cdr}(s)))))$,
 vs. $(\text{FORM}(s) \equiv (\text{ELF}(s) \vee \exists x \ f. ((s = x \ \text{gen} \ f) \vee (s = x \ \text{ex} \ f)) \vee \exists f1 \ f2. ((s = f1 \ \text{dis} \ f2) \vee (s = f1 \ \text{con} \ f2) \vee (s = f1 \ \text{impl} \ f2)) \vee \exists f. s = \text{neg}(f))) ; ;$

1.5 Sequences

AXIOM SEQ:

$\forall sq. \quad sq = \text{scar}(sq) \ c \ \text{scdr}(sq)$,
 $\forall sq1 \ sq2. \quad (sq1 = \text{SLAMBDA} \supset \text{scar}(sq1 \ c \ sq2) = \text{scar}(sq2))$,
 $\forall sq1 \ sq2. \quad (sq1 \neq \text{SLAMBDA} \supset \text{scar}(sq1 \ c \ sq2) = \text{scar}(sq1))$,
 $\forall sq1 \ sq2. \quad (sq1 = \text{SLAMBDA} \supset \text{scdr}(sq1 \ c \ sq2) = \text{scdr}(sq2))$,

$\forall sq1\ sq2. \quad (sq1 \neq SLAMBDA \supset scdr(sq1\ cc\ sq2) = scdr(sq1)\ cc\ sq2),$
 $vsq. \quad sq\ cc\ SLAMBDA = SLAMBDA\ cc\ sq,$
 $vsq. \quad sq\ cc\ SLAMBDA = sq,$
 $\forall sq1\ sq2\ sq3. \quad (sq1\ cc\ (sq2\ cc\ sq3) = (sq1\ cc\ sq2)\ cc\ sq3),$
 $vs. \quad (slen(s) = 1 \vee s = SLAMBDA),$
 $vsq. \quad slen(sq) \geq 0,$
 $\forall sq1\ sq2. \quad slen(sq1\ cc\ sq2) = slen(sq1) + slen(sq2),$
 $vsq. \quad 0\ sgl\ sq = SLAMBDA,$
 $vsq. \quad 1\ sgl\ sq = scar(sq),$
 $\forall n\ sq. \quad ((n > 1) \supset ((n\ sgl\ sq) = ((n-1)\ sgl\ scdr(sq)))) ;;$

AXIOM SUBSEQDEF:

$\forall n1\ n2\ sq1\ sq2. \quad (SUBSEP(sq1, sq2, n1, n2) \equiv (slen(sq2) = n2 - n1 + 1 \wedge$
 $(\forall n. (n \geq n1 \wedge n \leq n2 \supset n\ sgl\ sq2 = (n - n1 + 1)\ sgl\ sq1)))) ,$
 $\forall n1\ n2\ sq1\ sq2. \quad (SUBSEP(sq1, sq2, n1, n2) \equiv subseq(sq1, n1, n2) = sq2) ,$
 $\forall sq1\ sq2. \quad (SUBSSE(sq1, sq2) \equiv \exists n1\ n2. (SUBSEP(sq1, sq2, n1, n2))) ;;$

AXIOMEQSQ:

$\forall sq1\ sq2. \quad (\forall n. (n\ sgl\ sq1 = n\ sgl\ sq2) \supset sq1 = sq2) ;;$

1.6 Free and bound variables and the substitution

AXIOM BOUNDV:

$\forall x\ n\ f. \quad (GEB(x, n, f) \equiv \exists s1\ s2\ f1. (slen(s1) + 1 < n \wedge n < (slen(f) - slen(s2)) \wedge$
 $(x = n\ gl\ f) \wedge ((f = (s1\ c\ ((x\ gen\ f\ 1)\ c\ s2))) \vee (f = (s1\ c\ ((x\ ox\ f1)\ c\ s2))))) ;;$

AXIOM FREEV:

$\forall x\ n\ f. \quad (FRN(x, n, f) \equiv (x = (n\ gl\ f) \wedge \neg GEB(x, n, f))) ,$
 $\forall x\ f. \quad (FR(x, f) \equiv \exists n. FRN(x, n, f)) ;;$

AXIOM FIRSTFRDF:

$\forall x\ n\ f. \quad (FIRSTFREE(x, n, f) \equiv (FRN(x, n, f) \wedge \forall n1. (x = n1\ gl\ f \supset (n1 \geq n \vee GEB(x, n1, f))))) ,$
 $\forall x\ n\ f. \quad (FIRSTFREE(x, n, f) \equiv firstfreeocc(x, f) = n) ;;$

- AXIOM KFREEOCCDF:

$\forall x\ k\ n\ f. \quad (KTHFREEOCC(x, k, n, f) \equiv ((k = 0 \wedge n = 0) \vee$
 $(n = slen(f) \wedge \forall n2. (n2 > kthfreeocc(x, k-1, f) \supset \neg FRN(x, n2, f))) \vee$
 $(FRN(x, n, f) \wedge \forall n1. ((n1 < k \wedge n1 > 0) \supset \exists n2. (n2 < n \wedge KTHFREEOCC(x, n1, n2, f))))) ,$
 $\forall x\ k\ n\ f. \quad (KTHFREEOCC(x, k, n, f) \equiv kthfreeocc(x, k, f) = n) ,$
 $\forall x\ k\ n\ f. \quad (KTHFREEOCC(x, k, n, f) \supset numbfreeocc(x, n, f) = k),$
 $\forall x\ k\ n\ f. \quad (numbfreeocc(x, n, f) = k \supset (KTHFREEOCC(x, k, n, f) \vee$
 $(n < kthfreeocc(x, k, f) \wedge n > kthfreeocc(x, k-1, f)))) ;;$

AXIOM SUBSTDF:

$\forall x\ t\ f1\ f2. \quad (SBT(x, t, f1, f2) \equiv \forall n1\ n2. ((n2 = (numbfreeocc(x, n1, f1) * (slen(t) - 1)) + n1) \supset$
 $((\neg INDVAR(n1\ gl\ f1) \supset n1\ gl\ f1 = n2\ gl\ f2) \wedge$
 $(INDVAR(n1\ gl\ f1) \supset ((FRN(x, n1, f1) \supset SUBT(t, f2, n2)) \wedge$
 $(\neg FRN(x, n1, f1) \supset INVART(n1, f1, n2, f2)))))) ,$
 $\forall t\ f2\ n2. \quad (SUBT(t, f2, n2) \equiv \forall x2\ k. ((k\ gl\ t) = x2) \supset FRN(x2, n2 - (slen(t) - k), f2)) ,$
 $\forall n\ f1\ n1\ f2. \quad (INVART(n, f1, n1, f2) \equiv ((GEB(n1\ gl\ f2, n1, f2) \equiv GEB(n\ gl\ f1, n, f1)) \wedge$
 $(FRN(n1\ gl\ f2, n1, f2) \equiv FRN(n\ gl\ f1, n, f1)) \wedge n1\ gl\ f2 = n\ gl\ f1)) ,$
 $\forall x\ t\ f1\ f2. \quad (SBT(x, t, f1, f2) \equiv sbt(x, t, f1) = f2) ;;$

AXIOM SUBDEF:

$$\begin{aligned} & \forall x1 \ x2 \ f1 \ f2. (SBV(x1, x2, f1, f2) \equiv \forall n. ((\neg INDVAR(n \ gl \ f1) \supset n \ gl \ f1 = n \ gl \ f2) \wedge \\ & \quad (INDVAR(n \ gl \ f1) \supset ((FRN(x1, n, f1) \supset FRN(x2, n, f2)) \wedge \\ & \quad (\neg FRN(x1, n, f1) \supset INVARV(n, f1, f2)))))), \\ & \forall n \ f1 \ f2. (INVARV(n, f1, f2) \equiv ((GEB(n \ gl \ f2, n, f2) \equiv GEB(n \ gl \ f1, n, f1)) \wedge \\ & \quad (FRN(n \ gl \ f2, n, f2) \equiv FRN(n \ gl \ f1, n, f1)) \wedge n \ gt \ f2 = n \ gl \ f1)), \\ & \forall x \ x1 \ f1 \ f2. (SBV(x, x1, f1, f2) \equiv sbv(x, x1, f1) = f2)); \end{aligned}$$
1.7 Rules of inference**AXIOM ANDIRUL:**

$$\begin{aligned} & \text{Vsq pf1 pf2. } (ANDI(sq, pf1, pf2) \equiv \exists f1 \ f2. (scdr(sq) = (pf1 \ cc \ pf2) \wedge scar(sq) = f1 \ con \ f2 \wedge \\ & \quad f1 = scar(pf1) \wedge f2 = scar(pf2))), \\ & \text{vsq pf. } (ANDE(sq, pf) \equiv \exists f1 \ f2. (scdr(sq) = pf \wedge scar(sq) = f1 \wedge ((f1 \ con \ f2) = scar(pf)) \vee \\ & \quad (f2 \ con \ f1) = scar(pf)))); \end{aligned}$$
AXIOM FALSERUL:

$$\begin{aligned} & \text{Vsq pf1 pf2. } (FALSEI(sq, pf1, pf2) \equiv \exists f1. ((scdr(sq) = (pf1 \ cc \ pf2)) \wedge \\ & \quad (scar(sq) = FALSESYM) \wedge (\neg(f1) = scar(pf1)) \wedge (f1 = scar(pf2)))), \\ & \text{vsq pf. } (FALSEE(sq, pf) \equiv \exists f. ((scar(pf) = FALSESYM) \wedge f = scar(sq) \wedge scdr(sq) = pf)); \end{aligned}$$
AXIOM IMPLRUL:

$$\begin{aligned} & \text{Vsq pf1 pf2. } (IMPLE(sq, pf1, pf2) \equiv \exists f1 \ f2. ((scdr(sq) = (pf1 \ cc \ pf2)) \wedge \\ & \quad (scar(pf1) = (f1 \ impl \ f2)) \wedge (scar(sq) = f2 \wedge (scar(pf2) = f1))), \\ & \text{vsq pf f1. } (IMPLID(sq, pf, f1) \equiv (scdr(sq) = pf \wedge \exists f2. ((scar(sq) = (f1 \ impl \ f2)) \wedge \\ & \quad (f2 = scar(pf)) \wedge \exists n. (f1 = (n \ sgl \ pf))))), \\ & \text{vsq pf. } (IMPLI(sq, pf) \equiv \exists f. IMPLID(sq, pf, f)); \end{aligned}$$
AXIOM NEGRUL:

$$\begin{aligned} & \text{vsq pf f. } (NOTID(sq, pf, f) \equiv (scdr(sq) = pf \wedge scar(sq) = f \wedge (scar(pf) = FALSESYM) \wedge \\ & \quad \exists n. (n \ sgl \ pf) = f)), \\ & \text{vsq pf. } (NOTI(sq, pf) \equiv \exists f. NOTID(sq, pf, f)), \\ & \text{vsq pf f. } (NOTED(sq, pf, f) \equiv (scdr(sq) = pf \wedge (scar(pf) = FALSESYM) \wedge \\ & \quad \exists n. ((n \ sgl \ pf) = f) \wedge (f = \neg(scar(sq))))), \\ & \text{vsq pf. } (NOTE(sq, pf) \equiv \exists f. NOTED(sq, pf, f)); \end{aligned}$$
AXIOM ORRUL:

$$\begin{aligned} & \text{vsq pf. } (ORI(sq, pf) \equiv (scdr(sq) = pf \wedge \exists f1 \ f2. ((scar(sq) = (f1 \ dis \ f2)) \wedge \\ & \quad f1 = scar(pf) \vee f2 = scar(pf)))), \\ & \text{Vsq pf1 pf2 pf3 f1 f2. } (ORED(sq, pf1, pf2, pf3, f1, f2) \equiv (scdr(sq) = (pf1 \ cc \ (pf2 \ cc \ pf3)) \wedge \\ & \quad (scar(pf1) = (f1 \ dis \ f2) \wedge \exists f3. (scar(pf2) = f3 \wedge scar(sq) = f3 \wedge \\ & \quad (scar(pf3) = f3)) \wedge \exists n1. (n1 \ sgt \ pf2) = f1 \wedge \exists n1. (n1 \ sgt \ pf3) = f2))), \\ & \text{vsq pf1 pf2 pf3. } (ORE(sq, pf1, pf2, pf3) \equiv \exists f1 \ f2. ORED(sq, pf1, pf2, pf3, f1, f2)); \end{aligned}$$
AXIOM EXRUL:

$$\begin{aligned} & \text{vsq pf x t. } (EXI(sq, pf, x, t) \equiv \exists f1. ((scdr(sq) = pf1) \wedge (scar(sq) = (x \ ex \ f1)) \wedge \\ & \quad scar(pf) = sbt(x, t, f1))), \\ & \text{Vsq pf1 pf2 x1 x2 f1. } (EXED(sq, pf1, pf2, x1, x2, f1) \equiv ((scdr(sq) = (pf1 \ cc \ pf2)) \wedge \\ & \quad (scar(pf1) = (x1 \ ex \ f1)) \wedge (scar(sq) = scar(pf2)) \wedge \\ & \quad \exists n. ((n \ sgl \ pf2) = sbt(x1, x2, f1) \wedge EXAPPL(x2, pf2, f1)))), \\ & \text{vsq pf f1 pf2 x1 x2. } (EXAPPL(x, pf, f1) \equiv (\neg FR(x, scar(pf), D(sq, pf1, pf2, x1, x2, f1), \\ & \quad)) \wedge \neg FR(x, f1) \wedge \forall f1. (DEPEND(pf, f1) \supset \\ & \quad \neg FR(x, f1)))); \end{aligned}$$

AXIOM GENRUL:

$$\begin{aligned}
& \forall sq \, sq1 \, x \, t. \quad (GENE(sq, sq1, x, t) = (scdr(sq) = sq1 \wedge PROOFTREE(sq1) \wedge \\
& \quad \exists f. (scar(sq1) = x \text{ gen } f \wedge scar(sq) = sbt(x, t, f))))), \\
& \forall sq \, sq1 \, x1 \, x2. \quad (GENI(sq, sq1, x1, x2) = (scdr(sq) = sq1 \wedge PROOFTREE(sq1) \wedge \\
& \quad \exists f. (scar(sq) = x1 \text{ gen } f \wedge scar(sq1) = sbt(x1, x2, f) \wedge APGENI(x2, sq1))))), \\
& \forall x \, sq. \quad (APGENI(x, sq) = (\forall f. (DEPEND(sq, f) \supset \neg FR(x, f)) \wedge PROOFTREE(sq))), \\
& \forall pf. \exists x. \quad APGENI(x, pf);;
\end{aligned}$$

1.8 Deduction

AXIOM PROOF:

$$\begin{aligned}
& vsq. \quad (PROOFTREE(sq) = (FORM(sq) \vee \\
& \quad \exists pf. (ORI(sq, pf) \vee ANDE(sq, pf) \vee FALSEE(sq, pf) \vee NOTI(sq, pf) \vee NOTE(sq, pf) \vee \\
& \quad \quad IMPLI(sq, pf)) \vee \\
& \quad \exists pf \, x \, t. (GENI(sq, pf, x, t) \vee GENE(sq, pf, x, t) \vee EXI(sq, pf, x, t)) \vee \\
& \quad \exists pf1 \, pf2. (ANDI(sq, pf1, pf2) \vee FALSEI(sq, pf1, pf2) \vee IMPLI(sq, pf1, pf2)) \vee \\
& \quad \exists pf1 \, pf2 \, x1 \, x2. EXE(sq, pf1, pf2, x1, x2) \vee \\
& \quad \exists pf1 \, pf2 \, pf3. ORE(sq, pf1, pf2, pf3)));;
\end{aligned}$$

AXIOM DEPNDG:

$$\begin{aligned}
& vsq \, f. \quad (DEPEND(sq, f) \supset SUBSSE(f, sq)) , \\
& vsq \, f. \quad (f = sq \supset DEPEND(sq, f)) ;;
\end{aligned}$$

AXIOM DEPEND:

$$\begin{aligned}
& \forall pf \, pf1 \, pf2 \, f. \quad ((pf1 = scdr(pf) \supset (DEPEND(pf, f) = DEPEND(pf1, f))) = \\
& \quad (ORI(pf, pf1) \vee ANDE(pf, pf1) \vee FALSEE(pf, pf1) \vee \\
& \quad \exists f1. ((NOTID(pf, pf1, f1) \vee NOTED(pf, pf1, f1) \vee IMPLID(pf, pf1, f1)) \wedge f1 \neq f) \vee \\
& \quad \exists x \, t. (GENI(pf, pf1, x, t) \vee GENE(pf, pf1, x, t) \vee EXI(pf, pf1, x, t)))) , \\
& \forall pf \, pf1 \, pf2 \, f. \quad (((pf1 \text{ cc } pf2 = scdr(pf)) \vee (pf2 \text{ cc } pf1 = scdr(pf))) \supset \\
& \quad (DEPEND(pf, f) = ((DEPEND(pf1, f) \vee DEPEND(pf2, f)))) = \\
& \quad (ANDI(pf, pf1, pf2) \vee FALSEI(pf, pf1, pf2) \vee IMPLI(pf, pf1, pf2) \vee \\
& \quad \exists x1 \, x2 \, f1. (EXED(pf, pf1, pf2, x1, x2, f1) \wedge f \neq f1))), \\
& \forall pf \, pf1 \, pf2 \, pf3 \, f. \quad (((((pf1 \text{ cc } (pf2 \text{ cc } pf3)) = scdr(pf)) \vee \\
& \quad ((pf1 \text{ cc } (pf3 \text{ cc } pf2)) = scdr(pf)) \vee \\
& \quad ((pf2 \text{ cc } (pf1 \text{ cc } pf3)) = scdr(pf)) \vee \\
& \quad ((pf2 \text{ cc } (pf3 \text{ cc } pf1)) = scdr(pf)) \vee \\
& \quad ((pf3 \text{ cc } (pf1 \text{ cc } pf2)) = scdr(pf)) \vee \\
& \quad ((pf3 \text{ cc } (pf2 \text{ cc } pf1)) = scdr(pf))) \supset \\
& \quad (DEPEND(pf, f) = (DEPEND(pf1, f) \vee DEPEND(pf2, f) \vee \\
& \quad DEPEND(pf3, f)))) = \\
& \quad \exists f1 \, f2. (ORED(pf, pf1, pf2, pf3, f1, f2) \wedge f \neq f1 \wedge f \neq f2)) ;;
\end{aligned}$$

AXIOM NDEPND:

$$\begin{aligned}
& \forall pf1 \, pf2 \, f. \quad ((NOTID(pf1, pf2, f) \vee NOTED(pf1, pf2, f) \vee IMPLID(pf1, pf2, f)) \supset \\
& \quad \neg DEPEND(pf1, f)), \\
& \forall pf1 \, pf2 \, pf3 \, x1 \, x2 \, f1. \quad EXED(pf1, pf2, pf3, x1, x2, f1) \supset \neg DEPEND(pf1, f1) , \\
& \forall pf1 \, pf2 \, pf3 \, pf4 \, f1 \, f2. \quad (ORED(pf1, pf2, pf3, pf4, f1, f2) \supset DEPEND(pf1, f1) \wedge \\
& \quad \neg DEPEND(pf1, f2));; \\
& \forall f. \quad (BEW(f) = \exists sq. (PROOFTREE(sq) \wedge f = scar(sq) \wedge \forall f1. (DEPEND(sq, f1) \supset \\
& \quad AXIOM(f1)))));;
\end{aligned}$$

AXIOM THEORY:

$$\forall x \, f. \quad (AXIOM(f) \supset \neg FR(x, f));;$$

AXIOM INFVAR:

 $\forall s. \exists x. \forall n.$ $n \text{ gt } s \wedge x \text{ ;}$

APPENDIX 2

THE AXIOMS IN THE LOGIC

2.1 Natural numbers

AXIOM NUMB:

$$\begin{aligned}
& \forall n1 \ n2 \ n3. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2) \wedge \text{INTEGER}(n3)) \supset (n1=n2 \supset (n1=n3 \supset n2=n3))), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset (n1=n2 \supset \text{succ}(n1)=\text{succ}(n2))), \\
& \forall n1. (\text{INTEGER}(n1) \supset 0 \neq \text{succ}(n1)), \\
& \forall n1 \ n2. (\text{INTEGER}(n1) \wedge \text{INTEGER}(n2) \supset (\text{succ}(n1) = \text{succ}(n2) \supset n1 = n2)), \\
& \forall n1. (\text{INTEGER}(n1) \supset n1 + 0 = n1), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset n1 \cdot \text{succ}(n2) = \text{succ}(n1 + n2)), \\
& \forall n1. (\text{INTEGER}(n1) \supset n1 * 0 = 0), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset n1 * \text{succ}(n2) = (n1 * n2) + n1);;
\end{aligned}$$

AXIOM INDCT:

$$(F(0) \wedge \forall x. (\text{INTEGER}(x) \supset (F(x) \supset F(x+1)))) \supset \forall x. (\text{INTEGER}(x) \supset F(x));;$$

AXIOM DEFN:

$$\begin{aligned}
& \forall n. (\text{INTEGER}(n) \supset (\text{succ}(n)-1)=n), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset \text{succ}(n1)-n2=n1-(n2-1)), \\
& \forall n1 \ n2 \ n3. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2) \wedge \text{INTEGER}(n3)) \supset \\
& \quad (n1 < n2 \equiv \exists n3. (n3 \neq 0 \wedge n1 + n3 = n2))), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset (n1 \leq n2 \equiv (n1 < n2) \vee (n1 = n2))), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset (n2 > n1 \equiv n1 < n2)), \\
& \forall n1 \ n2. ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2)) \supset (n2 \geq n1 \equiv n1 \leq n2)),
\end{aligned}$$

2.2 The set of symbols

AXIOM SYM:

$$\begin{aligned}
& \forall s. (\text{SYM}(s) \equiv s = \text{LPARSYM} \vee s = \text{RPARSYM} \vee s = \text{ORSYM} \vee s = \text{ANDSYM} \vee s = \text{IMPSYM} \vee \\
& \quad s = \text{FALSESYM} \vee s = \text{FORALLSYM} \vee s = \text{EXISTSYM});;
\end{aligned}$$

2.3 Strings

AXIOM STRING:

$$\begin{aligned}
& \forall s. (\text{STRING}(s) \supset s = \text{car}(s) \text{ c } \text{cdr}(s)), \\
& \forall s1 \ s2. ((\text{STRING}(s1) \wedge \text{STRING}(s2)) \supset (s1 = \text{LAMBDA} \supset \text{car}(s1 \text{ c } s2) = \text{car}(s2))), \\
& \forall s1 \ s2. ((\text{STRING}(s1) \wedge \text{STRING}(s2)) \supset (s1 \neq \text{LAMBDA} \supset \text{car}(s1 \text{ c } s2) = \text{car}(s1))), \\
& \forall s1 \ s2. ((\text{STRING}(s1) \wedge \text{STRING}(s2)) \supset (s1 = \text{LAMBDA} \supset \text{cdr}(s1 \text{ c } s2) = \text{cdr}(s2))), \\
& \forall s1 \ s2. ((\text{STRING}(s1) \wedge \text{STRING}(s2)) \supset (s1 \neq \text{LAMBDA} \supset \text{cdr}(s1 \text{ c } s2) = \text{cdr}(s1))), \\
& \forall s. ((\text{STRING}(s) \supset (s \text{ c } \text{LAMBDA} = \text{LAMBDA} \text{ c } s)), \\
& \forall s. (\text{STRING}(s) \supset (s \text{ c } \text{LAMBDA} = s)), \\
& \forall s1 \ s2 \ s3. ((\text{STRING}(s1) \wedge \text{STRING}(s2) \wedge \text{STRING}(s3)) \supset (s1 \text{ c } (s2 \text{ c } s3) = (s1 \text{ c } s2) \text{ c } s3)), \\
& \forall s. (\text{STRING}(s) \supset (\text{len}(s) = 1 \vee s = \text{LAMBDA})), \\
& \forall s. (\text{STRING}(s) \supset \text{len}(s) \geq 0), \\
& \forall s1 \ s2. ((\text{STRING}(s1) \wedge \text{STRING}(s2)) \supset \text{len}(s1 \text{ c } s2) = \text{len}(s1) + \text{len}(s2)), \\
& \forall s. (\text{STRING}(s) \supset (\text{len}(s) = 1 \supset \text{ATOM}(s))),
\end{aligned}$$

$\forall s.$ (STRING(s) $\supset$ $\emptyset$ gl s=LAMBDA),
 $\forall s.$ (STRING(s) $\supset$ 1 gl s=car(s)),
 $\forall s \ n.$ ((STRING(s) $\wedge$ INTEGER(n)) $\supset$ ((n>1) $\supset$ ((n gl s)=((n-1) gl cdr(s))))),

AXIOM SUBSTRDEF:

$\forall n1 \ n2 \ s1 \ s2.$ ((INTEGER(n1) $\wedge$ INTEGER(n2) $\wedge$ STRING(s1) $\wedge$ STRING(s2)) $\supset$
 (SUBSTP(s1,s2,n1,n2) $\equiv$ (len(s2)=n2-n1+1 $\wedge$ ($\forall n.(n \geq n1 \wedge n \leq n2 \supset$
 $n \text{ gl } s1 = (n-n1+1) \text{ gl } s2$))))),
 $\forall n1 \ n2 \ s1 \ s2.$ ((INTEGER(n1) $\wedge$ INTEGER(n2) $\wedge$ STRING(s1) $\wedge$ STRING(s2)) $\supset$
 (SUBSTP(s1,s2,n1,n2) $\equiv$ substring(s1,n1,n2)=s),
 $\forall s1 \ 02.$ ((STRING(s1) $\wedge$ STRING(s2)) $\supset$ (SUBS(s1,s2) = $\exists n1 \ n2.$ SUBSTP(s1,s2,n1,n2)));

AXIOM DISEQ:

$\forall g1 \ g2.$ ($\neg(g1=g2) = g1 \neq g2$);;

AXIOM EQS:

$\forall s1 \ s2.$ ((STRING(s1) $\wedge$ STRING(s2)) $\supset$ ($\forall n.$ (INTEGER(n) $\supset$ (n gl s1 = n gl s2)) = s1 = s2)));;

AXIOM COMP:

$\forall f.$ (FORM(f) $\supset$ (e(f)=(LPARSYM c f) c RPARSYM)),
 $\forall f1 \ f2.$ ((FORM(f1) $\wedge$ FORM(f2)) $\supset$ (f1 dis f2)=(e(f1) c ORSYM) c e(f2)),
 $\forall f1 \ f2.$ ((FORM(f1) $\wedge$ FORM(f2)) $\supset$ (f1 impl f2)=(e(f1) c IMPSYM) c e(f2)),
 $\forall f.$ (FORM(f) $\supset$ neg(f)=(f impl FALSESYM)),
 $\forall f1 \ f2.$ ((FORM(f1) $\wedge$ FORM(f2)) $\supset$ (f1 con f2)=(e(f1) c ANDSYM) c e(f2)),
 $\forall x \ f2.$ ((INDVAR(x) $\wedge$ FORM(f2)) $\supset$ (x gen f2)=(FORALLSYM c x) c f2),
 $\forall x \ f2.$ ((INDVAR(x) $\wedge$ FORM(f2)) $\supset$ (x o x f2)=(EXISTSYM c x) c f2) ;;

2.4 Sequences

AXIOM SEQ:

$\forall sq.$ (SEQUENCE(sq) $\supset$ sq=scar(sq) cc scdr(sq)),
 $\forall sq1 \ sq2.$ ((SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2)) $\supset$ (sq1 =SLAMBDA $\supset$
 scar(sq1 cc sq2)=scar(sq2))),
 $\forall sq1 \ sq2.$ ((SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2)) $\supset$ (sq1 $\neq$ SLAMBDA $\supset$
 scar(sq1 cc sq2)=scar(sq1))),
 $\forall sq1 \ sq2.$ ((SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2)) $\supset$ (sq1 =SLAMBDA $\supset$
 scdr(sq1 cc sq2)=scdr(sq2))),
 $\forall sq1 \ sq2.$ ((SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2)) $\supset$ (sq1 $\neq$ SLAMBDA $\supset$
 scdr(sq1 cc sq2)=scdr(sq1) cc sq2)),
 $\forall sq.$ (SEQUENCE(sq) $\supset$ sq cc SLAMBDA=SLAMBDA cc sq),
 $\forall sq.$ (SEQUENCE(sq) $\supset$ sq cc SLAMBDA=sq),
 $\forall sq1 \ sq2 \ sq3.$ ((SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2) $\wedge$ SEQUENCE(sq3)) $\supset$
 (sq1 cc (sq2 cc sq3)=(sq1 cc sq2) cc sq3)),
 $\forall s.$ (STRING(s) $\supset$ (slen(s)= 1 $\vee$ s=SLAMBDA)),
 $\forall sq.$ (SEQUENCE(sq) $\supset$ slen(sq) $\geq$ 0),
 $\forall sq1 \ sq2.$ ((SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2)) $\supset$ slen(sq1 cc sq2)=slen(sq1)+slen(sq2)),
 $\forall sq.$ (SEQUENCE(sq) $\supset$ 8 sgl sq=SLAMBDA),
 $\forall sq.$ (SEQUENCE(sq) $\supset$ 1 sgl sq=scar(sq)),
 $\forall n \ sq.$ ((INTEGER(n) $\wedge$ SEQUENCE(sq)) $\supset$ ((n>1) $\supset$ ((n sgl sq)=((n-1) sgl scdr(sq)))));;

AXIOM SUBSEQDEF:

$\forall n1 \ n2 \ sq1 \ sq2.$ ((INTEGER(n1) $\wedge$ INTEGER(n2) $\wedge$ SEQUENCE(sq1) $\wedge$ SEQUENCE(sq2)) $\supset$

$$\begin{aligned}
& (\text{SUBSEP}(sq1, sq2, n1, n2) \equiv (\text{slen}(sq2) = n2 - n1 \cdot 1 \wedge (\forall n. (n \geq n1 \wedge n \leq n2 \Rightarrow \\
& \quad n \text{ sgl } sq2 = (n - n1 \cdot 1) \text{ sgl } sq1))))), \\
\forall n1 \ n2 \ sq1 \ sq2. & ((\text{INTEGER}(n1) \wedge \text{INTEGER}(n2) \wedge \text{SEQUENCE}(sq1) \wedge \text{SEQUENCE}(sq2)) \Rightarrow \\
& (\text{SUBSEP}(sq1, sq2, n1, n2) \equiv \text{subseq}(sq1, n1, n2) = sq2)), \\
\forall sq1 \ sq2. & ((\text{SEQUENCE}(sq1) \wedge \text{SEQUENCE}(sq2)) \Rightarrow (\text{SUBSSE}(sq1, sq2) = \\
& \exists n1 \ n2. (\text{SUBSEP}(sq1, sq2, n1, n2)))));;
\end{aligned}$$

AXIOM EQSQ:
 $\forall sq1 \ sq2. ((\text{SEQUENCE}(sq1) \wedge \text{SEQUENCE}(sq2)) \Rightarrow (\forall n. (n \text{ sgl } sq1 = n \text{ sgl } sq2) \Rightarrow sq1 = sq2));;$

2.5 Formulas

AXIOM FIND:
 $\begin{aligned} & \text{vsq.} \quad (\text{FIND}(\emptyset, \text{LAMBDA}, sq) \equiv \text{SEQUENCE}(sq)), \\ & \forall n \ s \ sq. \quad (\text{FIND}(n, s, sq) \equiv \text{INTEGER}(n) \wedge \text{STRING}(s) \wedge \text{SEQUENCE}(sq) \wedge \\ & \quad \exists n1 \ s1 \ s2. (\text{INTEGER}(n) \wedge \text{STRING}(s1) \wedge \text{STRING}(s2) \wedge (\emptyset < s \wedge s < \text{slen}(sq)) \wedge \\ & \quad (s1 = (n \text{ sgl } sq)) \wedge (s = (s1 \text{ c } s2)) \wedge \text{FIND}(n-1, s2, sq))));; \end{aligned}$

AXIOM FINDTOP:
 $\begin{aligned} & \text{vsq.} \quad (\text{FINDTOP}(\emptyset, \text{SLAMBDA}, sq) \equiv \text{SEQUENCE}(sq)), \\ & \forall n \ s \ sq. \quad (\text{FINDTOP}(n, s, sq) \equiv \text{INTEGER}(n) \wedge \text{STRING}(s) \wedge \text{SEQUENCE}(sq) \wedge \\ & \quad \exists s1 \ s2. (\text{STRING}(s1) \wedge \text{STRING}(s2) \wedge (s1 \neq \text{LAMBDA}) \wedge (s = (s1 \text{ c } s2)) \wedge \\ & \quad (s = \text{scar}(sq)) \wedge \text{FINDTOP}(n-1, s2, \text{scar}(sq))));; \end{aligned}$

AXIOM TERM:
 $\begin{aligned} & \text{vsq.} \quad (\text{TERMSEQ}(sq) \equiv \text{SEQUENCE}(sq) \wedge ((\text{slen}(sq) = 1 \wedge \text{INDVAR}(1 \text{ sgl } sq)) \vee \\ & \quad (\text{slen}(sq) > 1 \wedge \text{TERMSEQ}(\text{schr}(sq)) \wedge (\text{INDVAR}(\text{scar}(sq)) \vee \\ & \quad \exists n \ s. \text{INTEGER}(n) \wedge \text{STRING}(s) \wedge (s = \text{car}(\text{scar}(sq)) \wedge \text{OPCONST}(s) \wedge n = \text{arity}(s) \wedge \\ & \quad \text{FIND}(n, \text{cdr}(\text{scar}(sq)), \text{schr}(sq)))))), \\ & \forall t. \quad (\text{TERM}(t) \equiv \text{STRING}(t) \wedge \exists sq. (\text{TERMSEQ}(sq) \wedge t = \text{car}(sq)));; \end{aligned}$

AXIOM WFF:
 $\begin{aligned} & \forall f. \quad (\text{ELF}(f) \equiv \text{STRING}(f) \wedge (f = \text{FALSESYM} \vee \text{PREDPARO}(f) \vee \exists n \ sq. (\text{INTEGER}(n) \wedge \\ & \quad \text{SEQUENCE}(sq) \wedge \text{PREDPAR}(\text{car}(f)) \wedge n = \text{arity}(\text{car}(f)) \wedge \text{TERMSEQ}(sq) \wedge \\ & \quad \text{FINDTOP}(n, \text{cdr}(f), sq))), \\ & \text{vsq.} \quad (\text{FRR}(sq) \equiv \text{SEQUENCE}(sq) \wedge (sq \neq \text{SLAMBDA}) \wedge (\text{ELF}(\text{scar}(sq)) \vee \\ & \quad (\text{FRR}(\text{schr}(sq)) \wedge \exists s1 \ s2. (\text{STRING}(s1) \wedge \text{STRING}(s2) \wedge \\ & \quad (((\text{scar}(sq) = \text{neg}(s1) \wedge \text{FIND}(1, s1, \text{schr}(sq))) \vee \\ & \quad (\text{scar}(sq) = (s1 \text{ dis } s2) \wedge \text{FIND}(2, (s1 \text{ c } s2), \text{schr}(sq))) \vee \\ & \quad (\text{scar}(sq) = (s1 \text{ con } s2) \wedge \text{FIND}(2, (s1 \text{ c } s2), \text{schr}(sq))) \vee \\ & \quad (\text{scar}(sq) = (s1 \text{ impl } s2) \wedge \text{FIND}(2, (s1 \text{ c } s2), \text{schr}(sq))) \vee \\ & \quad (\text{scar}(sq) = (s1 \text{ gen } s2) \wedge \text{INDVAR}(s1) \wedge \text{FIND}(1, s2, \text{schr}(sq))) \vee \\ & \quad (\text{scar}(sq) = (s1 \text{ ex } s2) \wedge \text{INDVAR}(s1) \wedge \text{FIND}(1, s2, \text{schr}(sq))))))), \\ & \forall f. \quad (\text{FORM}(f) \equiv \text{STRING}(f) \wedge \exists sq. (\text{FRR}(sq) \wedge f = \text{scar}(sq)));; \end{aligned}$

2.6 Free and bound variables and the substitution

AXIOM BOUNDV:
 $\begin{aligned} & \forall x \ n \ f. (\text{GEB}(x, n, f) \equiv \text{INDVAR}(x) \wedge \text{INTEGER}(n) \wedge \text{FORM}(f) \wedge \exists s1 \ s2 \ f1. (\text{STRING}(s1) \wedge \\ & \quad \text{FORM}(f1) \wedge \text{STRING}(s2) \wedge \text{len}(s1) \cdot 1 < n \wedge n < (\text{len}(f) - \text{len}(s2)) \wedge \\ & \quad (x = n \text{ gl } f) \wedge ((f = (s1 \text{ c } ((x \text{ gen } f1) \text{ c } s2))) \vee (f = (s1 \text{ c } ((x \text{ ex } f1) \text{ c } f3))))));; \end{aligned}$

AXIOM FREEV:

$$\begin{aligned} \forall x \ n \ f. \quad & (FRN(x,n,f) \equiv INDVAR(x) \wedge INTEGER(n) \wedge FORM(f) \wedge x = (n \text{ gl } f) \wedge \\ & \neg GEB(x,n,f)), \\ \forall x \ f. \quad & (FR(x,f) \equiv \exists n. (INTEGER(n) \wedge FRN(x,n,f))) ;; \end{aligned}$$

AXIOM FIRSTFRDF:

$$\begin{aligned} \forall x \ n \ f. \quad & (FIRSTFREE(x,n,f) \equiv FRN(x,n,f) \wedge \forall n1. (INTEGER(n1) \wedge x = n1 \text{ gl } f \supset \\ & (n1 \geq n \vee GEB(x,n1,f))))), \\ \forall x \ n \ f. \quad & (FIRSTFREE(x,n,a) \equiv firstfree(x,f)=n) ;; \end{aligned}$$

AXIOM KFREEOCCDF:

$$\begin{aligned} \forall x \ k \ n \ f. \quad & (KTHFREEOCC(x,k,n,f) \equiv (INDVAR(x) \wedge INTEGER(k) \wedge INTEGER(n) \wedge \\ & FORM(f) \wedge (k=B \wedge n=B) \vee \\ & (n=len(f) \wedge \forall n2. ((INTEGER(n2) \wedge n2 > kthfreeocc(x,k-1,f)) \supset \neg FRN(x,n2,f))) \vee \\ & (FRN(x,n,f) \wedge \forall n1. ((INTEGER(n1) \wedge (n1 < k \wedge n1 > B)) \supset \\ & \exists n2. (INTEGER(n2) \wedge n2 < n \wedge KTHFREEOCC(x,n1,n2,f)))))), \\ \forall x \ k \ n \ f. \quad & (KTHFREEOCC(x,k,n,f) \equiv kthfreeocc(x,k,f)=n), \\ \forall x \ k \ n \ f. \quad & (KTHFREEOCC(x,k,n,f) \supset numbfreeocc(x,n,f)=k), \\ \forall x \ k \ n \ f. \quad & (numbfreeocc(x,n,f)=k \supset (KTHFREEOCC(x,k,n,f) \vee \\ & (n < kthfreeocc(x,k,f) \wedge n > kthfreeocc(x,k-1,f)))) ;; \end{aligned}$$

AXIOM SUBSTDF:

$$\begin{aligned} \forall x \ t \ f1 \ f2. \quad & (SBT(x,t,f1,f2) \equiv ((INDVAR(x) \wedge TERM(t) \wedge FORM(f1) \wedge FORM(f2)) \supset \\ & \forall n1 \ n2. ((INTEGER(n1) \wedge INTEGER(n2) \wedge \\ & n2 = numbfreeocc(x,n1,f1) * (len(t)-1) + n1 \supset \\ & ((\neg INDVAR(n1 \text{ gl } f1) \supset n1 \text{ gl } f1 = n2 \text{ gl } f2) \wedge \\ & (INDVAR(n1 \text{ gl } f1) \supset ((FRN(x,n1,f1) \supset SUBT(t,f2,n2)) \wedge \\ & (\neg FRN(x,n1,f1) \supset INVART(n1,f1,n2,f2))))))), \\ \forall t \ f2 \ n2. \quad & (SUBT(t,f2,n2) \equiv (TERM(t) \wedge FORM(f2) \wedge INTEGER(n2) \wedge \\ & \forall x2 \ k. ((INDVAR(x2) \wedge INTEGER(k) \wedge ((k \text{ gl } t) = x2)) \supset \\ & FRN(x2,n2-(len(t)-k),f2))))), \\ \forall n1 \ f1 \ n2 \ f2. \quad & (INVART(n1,f1,n2,f2) \equiv (INTEGER(n1) \wedge FORM(f1) \wedge INTEGER(n2) \wedge \\ & FORM(f2) \wedge (GEB(n2 \text{ gl } f2,n2,f2) \equiv GEB(n1 \text{ gl } f1,n1,f1)) \wedge \\ & (FRN(n2 \text{ gl } f2,n2,f2) \equiv FRN(n1 \text{ gl } f1,n1,f1)) \wedge \\ & n2 \text{ gl } f2 = n1 \text{ gl } f1)), \\ \forall x \ t \ f1 \ f2. \quad & ((INDVAR(x) \wedge TERM(t) \wedge FORM(f1) \wedge FORM(f2)) \supset \\ & (SBT(x,t,f1,f2) \equiv sbl(x,t,f1)=f2)), \\ \forall x \ t \ f1. \quad & ((INDVAR(x) \wedge TERM(t) \wedge FORM(f1)) \supset FORM(sbl(x,t,f1))) ;; \end{aligned}$$

AXIOM SUBDEF:

$$\begin{aligned} \forall x1 \ x2 \ f1 \ f2. \quad & (SBV(x1,x2,f1,f2) \equiv ((INDVAR(x1) \wedge INDVAR(x2) \wedge FORM(f1) \wedge FORM(f2)) \supset \\ & \forall n. (INTEGER(n) \supset ((\neg INDVAR(n \text{ gl } f1) \supset n \text{ gl } f1 = n \text{ gl } f2) \wedge \\ & (INDVAR(n \text{ gl } f1) \supset ((FRN(x1,n,f1) \supset FRN(x2,n,f2)) \wedge \\ & (\neg FRN(x1,n,f1) \supset INVARTV(n,f1,f2))))))), \\ \forall n \ f1 \ f2. \quad & (INVARTV(n,f1,f2) \equiv (INTEGER(n) \wedge FORM(f1) \wedge FORM(f2) \wedge \\ & (GEB(n \text{ gl } f2,n,f2) \equiv GEB(n \text{ gl } f1,n,f1)) \wedge \\ & FRN(n \text{ gl } f2,n,f2) \equiv FRN(n \text{ gl } f1,n,f1)) \wedge \\ & n \text{ gl } f2 = n \text{ gl } f1)), \\ \forall x1 \ x2 \ f1 \ f2. \quad & ((INDVAR(x1) \wedge INDVAR(x2) \wedge FORM(f1) \wedge FORM(f2)) \supset \\ & (SBV(x1,x2,f1,f2) \equiv sbv(x1,x2,f1)=f2)), \\ \forall x1 \ x2 \ f1. \quad & ((INDVAR(x1) \wedge INDVAR(x2) \wedge FORM(f1)) \supset FORM(sbv(x1,x2,f1))) ;; \end{aligned}$$

2.7 Rules of inference

AXIOM ANDIRUL:

vsq pfl pf2. $(\text{ANDI}(\text{sq}, \text{pf1}, \text{pf2}) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf1}) \text{ A } \text{PROOFTREE}(\text{pf2}) \text{ A } \exists f1 f2. (\text{schr}(\text{sq}) = (\text{pf1} \text{ cc } \text{pf2}) \text{ A } \text{scar}(\text{sq}) = f1 \text{ con } f2 \text{ A } \text{FORM}(f1) \text{ A } \text{FORM}(f2) \text{ A } f1 = \text{scar}(\text{pf1}) \text{ A } f2 = \text{scar}(\text{pf2}))))),$

vsq pf. $(\text{ANDE}(\text{sq}, \text{pf}) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE} \text{ A } \exists f1. (\text{schr}(\text{sq}) = \text{pf} \wedge \text{FORM}(f1) \text{ A } ((\text{scar}(\text{sq}) \text{ con } f1) = \text{scar}(\text{pf})) \vee ((f1 \text{ con } (\text{scar}(\text{sq})) = \text{scar}(\text{pf}))))))));$

AXIOM FALSERUL :

vsq pfl pf2. $(\text{FALSEI}(\text{sq}, \text{pf1}, \text{pf2}) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf1}) \text{ A } \text{PROOFTREE}(\text{pf2}) \text{ A } \exists f1. ((\text{schr}(\text{sq}) = (\text{pf1} \text{ cc } \text{pf2})) \text{ A } (\text{scar}(\text{sq}) = \text{FALSESYM}) \text{ A } \text{FORM}(f1) \text{ A } (\text{neg}(x) = \text{scar}(\text{pf1})) \text{ A } (x1 = \text{scar}(\text{pf2}))))),$

vsq pf. $(\text{FALSEE}(\text{sq}, \text{pf}) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf}) \text{ A } (\text{scar}(\text{pf}) = \text{FALSESYM}) \text{ A } \text{schr}(\text{sq}) = \text{pf}));$

AXIOM IMPLRUL :

Vsq pfl pf2. $(\text{IMPLE}(\text{sq}, \text{pf1}, \text{pf2}) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf1}) \text{ A } \text{PROOFTREE}(\text{pf2}) \text{ A } \forall f1. ((\text{schr}(\text{sq}) = (\text{pf1} \text{ cc } \text{pf2})) \text{ A } \text{FORM}(f1) \text{ A } (\text{scar}(\text{pf1}) = (f1 \text{ impl } (\text{scar}(\text{sq}))) \text{ A } (\text{scar}(\text{pf2}) = f1))))),$

vsq pf f1. $(\text{IMPLID}(\text{sq}, \text{pf}, f1) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf}) \text{ A } \text{schr}(\text{sq}) = \text{pf} \text{ A } \text{FORM}(f1) \text{ A } \exists f2. ((\text{scar}(\text{sq}) = (f1 \text{ impl } x2)) \text{ A } \text{FORM}(f1) \text{ A } (f2 = \text{scar}(\text{pf})) \wedge \exists n. (\text{INTEGER}(n) \text{ A } f1 = (n \text{ sgl } \text{pf}))))));$

vsq pf. $(\text{IMPLI}(\text{sq}, \text{pf}) \equiv \exists f. \text{IMPLID}(\text{sq}, \text{pf}, f));$

AXIOM NECRUL:

vsq pf f1. $(\text{NOTID}(\text{sq}, \text{pf}, f1) \equiv (\text{schr}(\text{sq}) = \text{pf} \text{ A } \text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf}) \text{ A } \text{FORM}(f1) \text{ A } \exists n. ((\text{scar}(\text{pf}) = \text{FALSESYM}) \text{ A } \text{scar}(\text{sq}) = \text{neg}(f1) \text{ A } \text{INTEGER}(n) \text{ A } ((n \text{ sgl } \text{pf}) = f1))))),$

vsq pf. $(\text{NOTI}(\text{sq}, \text{pf}) \equiv \exists f. \text{NOTID}(\text{sq}, \text{pf}, f)),$

vsq pf f1. $(\text{NOTED}(\text{sq}, \text{pf}, f1) \equiv (\text{schr}(\text{sq}) = \text{pf} \text{ A } \text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE} \text{ A } \text{FORM}(f) \text{ A } \exists n. ((\text{scar}(\text{pf}) = \text{FALSESYM}) \text{ A } \text{INTEGER}(n) \text{ A } ((n \text{ sgl } \text{pf}) = \text{neg}(\text{scar}(\text{sq}))))),$

vsq pf. $(\text{NOTE}(\text{sq}, \text{pf}) \equiv \exists f. \text{NOTED}(\text{sq}, \text{pf}, f));$

AXIOM ORRUL:

vsq pf. $(\text{ORI}(\text{sq}, \text{pf}) \equiv (\text{schr}(\text{sq}) = \text{pf} \text{ A } \text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE} \text{ A } \exists f1 f2. ((\text{scar}(\text{sq}) = (f1 \text{ dis } f2)) \text{ A } \text{FORM}(f1) \text{ A } \text{FORM}(f2) \text{ A } (f1 = \text{scar}(\text{pf})) \vee (f2 = \text{scar}(\text{pf}))))),$

Vsq pfl pf2 pf3 f1 f2. $(\text{ORED}(\text{sq}, \text{pf1}, \text{pf2}, \text{pf3}, f1, f2) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf1}) \text{ A } \text{PROOFTREE}(\text{pf2}) \text{ A } \text{PROOFTREE}(\text{pf3}) \text{ A } \text{FORM}(f1) \text{ A } \text{FORM}(f2) \text{ A } (\text{schr}(\text{sq}) = (\text{pf1} \text{ cc } (\text{pf2} \text{ cc } \text{pf3}))) \text{ A } (\text{scar}(\text{pf1}) = (f1 \text{ dis } f2)) \wedge (\text{scar}(\text{pf2}) = \text{scar}(\text{sq})) \text{ A } (\text{scar}(\text{pf3}) = \text{scar}(\text{sq})) \text{ A } \exists n1. (n1 \text{ sgl } \text{pf2}) = f1) \text{ A } \exists n1. (n1 \text{ sgl } \text{pf3}) = f2))))),$

Vsq pf1 pf2 pf3. $(\text{ORE}(\text{sq}, \text{pf1}, \text{pf2}, \text{pf3}) \equiv \exists f1 f2. \text{ORED}(\text{sq}, \text{pf1}, \text{pf2}, \text{pf3}, f1, f2));$

AXIOM EXRUL :

vsq pf x t. $(\text{EXI}(\text{sq}, \text{pf}, x, t) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf}) \text{ A } \text{INDVAR}(x) \text{ A } \text{TERM}(t) \wedge \exists f1. ((\text{schr}(\text{sq}) = \text{pf1}) \text{ A } (\text{scar}(\text{sq}) = (x \text{ ex } f1)) \text{ A } \text{FORM}(f1) \text{ A } \text{scar}(\text{pf}) = \text{sbt}(x, t, f1))))),$

Vsq pf1 pf2 x1 x2 f1. $(\text{EXED}(\text{sq}, \text{pf1}, \text{pf2}, x1, x2, f1) \equiv (\text{SEQUENCE}(\text{sq}) \text{ A } \text{PROOFTREE}(\text{pf1}) \text{ A } \text{INDVAR}(x1) \text{ A } \text{INDVAR}(x2) \text{ A } (\text{schr}(\text{sq}) = (\text{pf1} \text{ cc } \text{pf2})) \text{ A } \text{FORM}(f1) \text{ A } (\text{scar}(\text{pf1}) = (x1 \text{ ex } f1)) \text{ A } (\text{scar}(\text{sq}) = \text{scar}(\text{pf2})) \text{ A } \exists n. ((n \text{ sgl } \text{pf2}) = \text{sbt}(x1, x2, f1) \text{ A } \text{INTEGER}(n) \text{ A } \text{EXAPPL}(x2, \text{pf2}, f1))))),$

Vsq pf1 pf2 x1 x2. $(\text{EXE}(\text{sq}, \text{pf1}, \text{pf2}, x1, x2) \equiv \text{EXED}(\text{sq}, \text{pf1}, x1, x2)),$

vx pf f. $(\text{EXAPPL}(x, \text{pf}, f) \equiv (\text{INDVAR}(x) \text{ A } \text{PROOFTREE}(\text{pf}) \text{ A } \text{FORM}(f) \text{ A } \neg \text{FR}(x, \text{scar}(\text{pf})) \wedge$

$$\neg FR(x, f) \wedge \forall f. 1. (DEPEND(pf, f) \supset \neg FR(x, f));;$$

AXIOM GENRUL:

vsq sq1 x a. $(GENE(sq, sq1, x, t) \equiv (SEQUENCE(sq) \wedge INDVAR(x) \wedge TERM(t) \wedge scdr(sq) = sq1 \wedge PROOFTREE(sq1) \wedge \exists f. (FORM(f) \wedge scar(sq1) = x \text{ gen } f \wedge scar(sq) = sbt(x, t, f))));$

vsq sq1 x1 x2. $(GENI(sq, sq1, x1, x2) \equiv (SEQUENCE(sq) \wedge INDVAR(x1) \wedge INDVAR(x2) \wedge scdr(sq) = sq1 \wedge PROOFTREE(sq1) \wedge \exists f. (FORM(f) \wedge (scar(sq) = x1 \text{ gen } f) \wedge scar(sq1) = sbt(x1, x2, f) \wedge APGENI(x2, sq1))));$

vx sq. $(APGENI(x, sq) \equiv (INDVAR(x) \wedge \forall f. (DEPEND(sq, f) \supset \neg FR(x, f))) \wedge PROOFTREE(sq));$

vsq. $(PROOFTREE(sq) \supset \exists x. (INDVAR(x) \wedge APGENI(x, sq))) ;;$

2.8 Deduction

AXIOM PROOF:

$\forall sq.$ $(PROOFTREE(sq) \equiv ((SEQUENCE(sq) \wedge FORM(sq)) \vee \exists pf. (PROOFTREE(pf) \wedge (ORI(sq, pf) \vee ANDE(sq, pf) \vee FALSEE(sq, pf) \vee NOTI(sq, pf) \vee NOTE(sq, pf) \vee IMPLI(sq, pf))) \vee \exists pf \ x \ t. (PROOFTREE(pf) \wedge INDVAR(x) \wedge TERM(t) \wedge (GENI(sq, pf, x, t) \vee GENE(sq, pf, x, t) \vee EXI(sq, pf, x, t)))) \vee \exists pf \ 1 \ pf2. (PROOFTREE(pf1) \wedge PROOFTREE(pf2) \wedge (ANDI(sq, pf1, pf2) \vee FALSEI(sq, pf1, pf2) \vee IMPLI(sq, pf1, pf2))) \vee \exists pf \ 1 \ pf2 \ x1 \ x2. (PROOFTREE(pf1) \wedge PROOFTREE(pf2) \wedge INDVAR(x1) \wedge INDVAR(x2) \wedge EXE(sq, pf1, pf2, x1, x2)) \vee \exists pf \ 1 \ pf2 \ pf3. (PROOFTREE(pf1) \wedge PROOFTREE(pf2) \wedge PROOFTREE(pf3) \wedge ORE(sq, pf1, pf2, pf3)))) ;;$

AXIOM DEPNDG:

$\forall sq \ f.$ $(DEPEND(sq, f) \supset (SEQUENCE(sq) \wedge FORM(f) \wedge SUBSSE(f, sq))),$
 $\forall sq \ f. ((SEQUENCE(sq) \wedge FORM(f) \wedge sq = f) \supset DEPEND(sq, f)) ;;$

AXIOM DEPEND:

$\forall pf \ pf1 \ f.$ $((((PROOFTREE(pf) \wedge PROOFTREE(pf1) \wedge (pf1 = scdr(pf))) \supset (DEPEND(pf, f) \equiv DEPEND(pf1, f))) \equiv (ORI(pf, pf1) \vee ANDE(pf, pf1) \vee FALSEE(pf, pf1) \vee \exists f1. (FORM(f1) \wedge (NOTID(pf, pf1, f1) \vee NOTED(pf, pf1, f1) \vee IMPLID(pf, pf1, f1)) \wedge f1 \neq f) \vee \exists x \ t. (INDVAR(x) \wedge TERM(t) \wedge GENI(pf, pf1, x, t) \vee GENE(pf, pf1, x, t) \vee EXI(pf, pf1, x, t)))))) ;;$

AXIOM DEP:

$\forall pf \ pf1 \ pf2 \ f.$ $(((((PROOFTREE \wedge PROOFTREE(pf1) \wedge PROOFTREE(pf2) \wedge ((pf1 \text{ cc } pf2 = scdr(pf)) \vee (pf2 \text{ cc } pf1 = scdr(pf)))) \supset (DEPEND(pf, f) \equiv (DEPENI(pf1, f) \vee DEPEND(pf2, f)))) \equiv (ANDI(pf, pf1, pf2) \vee FALSEI(pf, pf1, pf2) \vee IMPLI(pf, pf1, pf2) \vee \exists x1 \ x2 \ f1. (EXED(pf, pf1, pf2, x1, x2, f1) \wedge f \neq f1)))) ;;$

AXIOM DEPND:

$\forall pf \ pf1 \ pf2 \ pf3 \ f.$ $(((((PROOFTREE \wedge PROOFTREE(pf1) \wedge PROOFTREE(pf2) \wedge PROOFTREE(pf3) \wedge (((pf1 \text{ cc } (pf2 \text{ cc } pf3)) = scdr(pf)) \vee ((pf1 \text{ cc } (pf3 \text{ cc } pf2)) = scdr(pf)) \vee$

$$\begin{aligned}
& ((pf2 \text{ cc } (pf1 \text{ cc } pf3)) = sdr(pf)) \vee \\
& ((pf2 \text{ cc } (pf3 \text{ cc } pf1)) = sdr(pf)) \vee \\
& ((pf3 \text{ cc } (pf1 \text{ cc } pf2)) = sdr(pf)) \vee \\
& ((pf3 \text{ cc } (pf2 \text{ cc } pf1)) = sdr(pf))) \supset \\
& (DEPEND(pf, f) \equiv (DEPEND(pf1, f) \vee DEPEND(pf2, f) \vee DEPEND(pf3, f))) \equiv \\
& \exists f1 f2. (ORED(pf, pf1, pf2, pf3, f1, f2) \wedge f \neq f1 \wedge f \neq f2) ,
\end{aligned}$$

AXIOM NDEPND:

$$\begin{aligned}
& \forall pf1 \text{ pf2 } f. ((NOTID(pf1, pf2, f) \vee NOTED(pf1, pf2, f) \vee tMPLtD(pf1, pf2, f)) \supset \\
& \quad \neg DEPEND(pf1, f)), \\
& \forall pf1 \text{ pf2 } pf3 \text{ xl } x2 \text{ f} (EXED(pf1, pf2, pf3, xl, x2, f) \supset \neg DEPEND(pf1, f)) , \\
& \forall pf1 \text{ pf2 } pf3 \text{ pf4 } f1 \text{ f2}. (ORED(pf1, pf2, pf3, pf4, f1, f2) \supset \neg DEPEND(pf1, f1) \wedge \neg DEPEND(pf1, f2));;
\end{aligned}$$

AXIOM PROVABLE:

$$\begin{aligned}
& \forall f. (BEW(f) \equiv FORM(f) \wedge \exists sq. (PROOFTREE(sq) \wedge f = scar(sq) \wedge \\
& \quad \forall f1. (DEPEND(sq, f1) \supset AXIOM(f1)))));;
\end{aligned}$$

AXIOM THEORY:

$$\forall x \text{ f}. (AXIOM(f) \supset \neg FR(x, f) \wedge FORM(f));;$$

AXIOM INFVAR:

$$\forall s. \exists x. \forall n. n \text{ gt } s \neq x);;$$

APPENDIX 3

THE PROOF OF "IF f IS A WFF ALSO $\lambda x.f$ IS A WFF"

3.1 FOL commands and printout in the many sorted logic commands

```

VE WFF1, x gen f;
TAUTEQ (x gen f = x gen f) v (x gen f = x ex f);
UNIFY --: #2 #1, -;
TAUT --: #1, 1 :-;

```

```

proof

```

```

1 FORM(x gen f) = (ELF(x gen f) v (exists f1. ((x gen f) = (x1 gen f 1) v (x gen f) = (x1 ex f1)) v
  (exists f2. ((x gen f) = (f1 dis f2) v ((x gen f) = (f1 con f2) v (x gen f) = (f1 impl f2))) v
  exists f1. (x gen f) = neg(f1))))

```

```

2 (x gen f) = (x gen f) v (x gen f) = (x ex f)

```

```

3 exists f1. ((x gen f) = (x1 gen f1) v (x gen f) = (x1 ex f1))

```

```

4 FORM(x gen f)

```

3.2 FOL commands in the earlier axiomatization

```

DECLARE INDVAR A U;
label hpt 1;
ASSUME FORM(f) A INDVAR (x1);
label teo1;
ASSUME V f s. (SEQUENCE(sq) ^ sq / SLAMBDA => (STRING(s) =, (s cc sq) / SLAMBDA));
label teo2;
ASSUME V s sq. (STRING(s) ^ SEQUENCE(sq) = scar(s cc sq) = s);
label teo3;
ASSUME V s sq. (STRING(s) ^ SEQUENCE(sq) = scdr(s cc sq) = sq);
label teo4;
ASSUME V sq. (SEQUENCE(sq) ^ sq / SLAMBDA = find(1, scar(sq), sq));
label teo5;
ASSUME V f x. (FORM(f) ^ INDVAR(x) => STRING(x gen f));
label teo6;
ASSUME V s sq. (STRING(s) ^ SEQUENCE(sq) => SEQUENCE(s cc sq));
label teo7;
ASSUME V x. (INDVAR(x) => STRING(x));

V e WFF2 f;
LABEL ass1;
taut exists sq. (FRR(sq) ^ f = scar(sq)) 1 :-;
ASSUME FRR(SQ) A f = SCAR(SQ);
V e WFF1 S Q;
V o teo1 SQ, x1 gen f;

```

```

Ve teo2 x1 gen f ,SQ;
Ve teo3 x1 gen f ,SQ;
Ve teo4      SQ;
Ve teo5 f ,x1;
Ve teo7 x1;
Ve Wff1 (x1 gen f) cc SQ;

TAUTEQ -:#2*2*2*2*2*1*1[s1←f : s2←x1] 1:-;
unify --:#2*2*2*2*2*2-;
Ve teo6 x1 gen f ,SQ;
Ve WFF2 x1 gen f;
tauteq -:#2*2*1[sq←(x1 gen f) cc SQ] 1:-;
unify --:#2*2*2-;
taut FORM(x1 gen f) 1:-;
Ze ass 1 ,-,SQ;
Di hpt1 ,-,;
Vi -,x1,f;

```

3.3 Printout of the proof in the earlier axiomatization

```

1 FORM(f)∧INDVAR(x1) (1) --- ASSUME

2 Vsq s.((SEQUENCE(sq)∧sq≠SLAMBDA)⇒(STRING(s)⇒(s cc sq)≠SLAMBDA)) (2) --- ASSUME

3 Vs sq.((STRING(s)∧SEQUENCE(sq))⇒scar(s cc sq)=s) (3) --- ASSUME

4 Vs sq.((STRING(s)∧SEQUENCE(sq))⇒scdr(s cc sq)=sq) (4) --- ASSUME

5 Vsq.((SEQUENCE(sq)∧sq≠SLAMBDA)⇒find(1,scar(sq),sq)) (5) --- ASSUME

6 Vf x.((FORM(f)∧INDVAR(x))⇒STRING(x gen f)) (6) --- ASSUME

7 Vs sq.((STRING(s)∧SEQUENCE(sq))⇒SEQUENCE(s cc sq)) (7) --- ASSUME

8 Vx.(INDVAR(x)⇒STRING(x)) (8) --- ASSUME

9 FORM(f)⇒(STRING(f)∧Essq.(FRR(sq)∧f=scar(sq))) --- VE WFF2 f

10 Essq.(FRR(sq)∧f=scar(sq)) (1 2 3 4 5 6 7 8) --- TAUT 1:9

1 1 FRR(SQ)∧f=scar(SQ) (1 1) --- ASSUME

1 2 FRR(SQ)⇒(SEQUENCE(SQ)∧(SQ≠SLAMBDA∧(ELF(scar(SQ))∨(FRR(scdr(SQ))∧Es1
s2.(STRING(s1)∧(STRING(s2)∧((scar(SQ)=NEG(s1)∧find(1,s1,scdr(SQ)))∨((scar(SQ)
=(s1 dis s2)∧find(2,s1 c x2,scdr(SQ)))∨((scar(SQ)=(s1 con s2)∧find(2,s1 c s2,
scdr(SQ)))∨((scar(SQ)=(s1 impl s2)∧find(2,s1 c s2,scdr(SQ)))∨((scar(SQ)=(s1 gen
s2)∧(INDVAR(s1)∧find(1,s2,scdr(SQ)))∨(scar(SQ)=(s1 ex s2)∧(INDVAR(s1)∧find(1,
s2,scdr(SQ)))))))))))))) - - VE WFF1 SQ

1 3 (SEQUENCE(SQ)∧SQ≠SLAMBDA)⇒(STRING(x1 gen f)⇒(x1 gen f) cc SQ)≠SLAMBDA (2)
--- VE 2 SQ,x1 gen f

```

14 $(\text{STRING}(x1 \text{ gen } f) \wedge \text{SEQUENCE}(SQ)) = \text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (x1 \text{ gen } f)$
 (3) --- VE 3 x1 gen f, SQ
 15 $(\text{STRING}(x1 \text{ gen } f) \wedge \text{SEQUENCE}(SQ)) = \text{schr}((x1 \text{ gen } f) \text{ cc } SQ) = SQ$ (4) --- VE 4 x1 gen f, SQ
 16 $(\text{SEQUENCE}(SQ) \wedge SQ / \text{SLAMBDA}) = \text{find}(1, \text{scar}(SQ), SQ)$ (5) --- VE 5 SQ
 17 $(\text{FORM}(f) \wedge \text{INDVAR}(x1)) \supset \text{string}(x1 \text{ gen } f)$ (6) --- VE 6 f, x1
 18 $\text{INDVAR}(x1) \supset \text{STRING}(x1)$ (8) --- VE 8 x1
 19 $\text{FRR}((x1 \text{ gen } f) \text{ cc } SQ) = (\text{SEQUENCE}((x1 \text{ gen } f) \text{ cc } SQ) \wedge ((x1 \text{ gen } f) \text{ cc } U) / \text{SLAMBDA} \wedge (\text{ELF}(\text{scar}((x1 \text{ gen } f) \text{ cc } SQ)) \vee (\text{FRR}(\text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \wedge \exists s1 s2. (\text{STRING}(s1) \wedge (\text{STRING}(s2) \wedge ((\text{scar}((s1 \text{ gen } f) \text{ cc } SQ) = \text{NEG}(s1) \wedge \text{find}(1, s1, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ dis } s2) \wedge \text{find}(2, s1 \text{ c } s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ con } s2) \wedge \text{find}(2, s1 \text{ c } s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ impl } s2) \wedge \text{find}(2, s1 \text{ c } s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ gen } s2) \wedge (\text{INDVAR}(s1) \wedge \text{find}(1, s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee (\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ ex } s2) \wedge (\text{INDVAR}(s1) \wedge \text{find}(1, s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)))))))))))))$ --- VE WFF1 (x1 gen f) c c SQ
 20 $\text{STRING}(x1) \wedge (\text{STRING}(f) \wedge ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = \text{NEG}(x1) \wedge \text{find}(1, x1, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (x1 \text{ dis } f) \wedge \text{find}(2, x1 \text{ c } f, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (x1 \text{ con } f) \wedge \text{find}(2, x1 \text{ c } f, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (x1 \text{ impl } f) \wedge \text{find}(2, x1 \text{ c } f, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (x1 \text{ gen } f) \wedge (\text{INDVAR}(x1) \wedge \text{find}(1, f, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee (\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (x1 \text{ ex } f) \wedge (\text{INDVAR}(x1) \wedge \text{find}(1, f, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)))))))))))))$ (1 2 3 4 5 6 7 8 11) --- TAUTEQ 1:19
 21 $\exists s1 s2. (\text{STRING}(s1) \wedge (\text{STRING}(s2) \wedge ((\text{scar}((s1 \text{ gen } f) \text{ c c } SQ) = \text{NEG}(s1) \wedge \text{find}(1, s1, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ dis } s2) \wedge \text{find}(2, s1 \text{ c } s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ con } s2) \wedge \text{find}(2, s1 \text{ c } s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ impl } s2) \wedge \text{find}(2, s1 \text{ c } s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee ((\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ gen } s2) \wedge (\text{INDVAR}(s1) \wedge \text{find}(1, s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)) \vee (\text{scar}((x1 \text{ gen } f) \text{ cc } SQ) = (s1 \text{ ex } s2) \wedge (\text{INDVAR}(s1) \wedge \text{find}(1, s2, \text{schr}((x1 \text{ gen } f) \text{ cc } SQ)))))))))))))$ (1 2 3 4 5 6 7 8 11) --- UNIFY 20
 22 $(\text{STRING}(x1 \text{ gen } f) \wedge \text{SEQUENCE}(SQ)) \supset \text{SEQUENCE}((x1 \text{ gen } f) \text{ cc } SQ)$ (7) --- VE 7 x1 GEN f, SQ
 23 $\text{FORM}(x1 \text{ gen } f) = (\text{STRING}(x1 \text{ gen } f) \wedge \exists sq. (\text{FRR}(sq) \wedge (x1 \text{ gen } f) = \text{scar}(sq)))$ --- VE WFF2 x1 gen f
 24 $\text{FRR}((x1 \text{ gen } f) \text{ cc } SQ) \wedge (x1 \text{ gen } f) = \text{scar}((x1 \text{ gen } f) \text{ cc } SQ)$ (1 2 3 4 5 6 7 8 11) TAUTEQ 1:23
 25 $\exists sq. (\text{FRR}(sq) \wedge (x1 \text{ gen } f) = \text{scar}(sq))$ (1 2 3 4 5 6 7 8 11) --- UNIFY 24
 26 $\text{FORM}(x1 \text{ gen } f)$ (1 2 3 4 5 6 7 8 11) --- TAUT 1:25
 27 $\text{FORM}(x1 \text{ gen } f)$ (1 2 3 4 5 6 7 8) --- $\exists E$ 10 26 U
 28 $(\text{FORM}(f) \wedge \text{INDVAR}(x1)) \supset \text{FORM}(x1 \text{ gen } f)$ (2 3 4 5 6 7 8) --- $\supset I$ 2 7
 29 $\forall f x1. ((\text{FORM}(f) \wedge \text{INDVAR}(x1)) \supset \text{FORM}(x1 \text{ gen } f))$ (2 3 4 5 6 7 8) --- $\forall I$ 28 x1 $\leftarrow$ x1 f $\leftarrow$ f

APPENDIX 4

THE PROOF OF THE EQUIVALENCE BETWEEN **SBV** AND **SBT** FOR VARIABLES

4.1 FOL commands in the many sorted logic

```

LABEL ARITH1 ; ASSUME  $\forall n. (n * (\text{len}(x) - 1) = 0)$ ;
LABEL ARITH2; ASSUME  $\forall n. (0 + n = n)$ ;
LABEL ARITH3; ASSUME  $\forall x. (\text{len}(x) - 1) = 0$ ;
LABEL ARITH4; ASSUME  $\forall n. (n - 0) = n$ ;
LABEL STRING1 ; ASSUME  $\forall x. \text{len } x = x$ ;

```

Proof of the First Lemma: $\forall x \ f \ n. (\text{SUBT}(x, f, n) \supset \text{FRN}(x, n, f))$

```

LABEL HPTLEM; ASSUME SUBT(x, f, n);
 $\forall e$  SUBSTDF 1, x, f, n;
TAUT  $\neg : \#2, \neg, \neg$ ;
 $\forall e \neg, x, 1$ ;
 $\forall e$  STRING1, x; substr = in  $\neg$ ;
 $\forall e$  ARITH1, x; substr = in  $\neg$ ;
 $\forall e$  ARITH4, n; substr = in  $\neg$ ;
TAUTEQ FRN(x, n, f), HPTLEM + 1;  $\neg$ ;
 $\supset$ ! HPTLEM,  $\neg$ ;
LABEL LEMMA 1 ;  $\forall i \neg, x, f, n$ ;

```

Proof of the Second Lemma: $\forall n \ f \ 1 \ f2. (\text{INVART}(n, f \ 1 \ , n, f2) = \text{INVARV}(n, f1, f2))$

```

 $\forall e$  SUBSTDF2, n, f 1, n, f2;
 $\forall e$  SUBDEF 1, n, f 1, f2;
TAUT  $\neg : \#1 = \neg : \#1, \neg, \neg$ ;
LABEL LEMMA2;  $\forall i \neg, n, f1, f2$ ;

```

Proof of the Main Theorem: $\forall x1 \ x2 \ f \ 1 \ f2. (\text{SBT}(x1, x2, f \ 1 \ , f2) \supset \text{SBV}(x1, x2, f1, f2))$

```

LABEL HPT; ASSUME SBT(x1, x2, f1, f2);
 $\forall e$  SUBSTDF0, x1, x2, f1, f2;
TAUT  $\neg : \#2, \text{HPT}, \neg$ ;
 $\forall e \neg, n1, n1$ ;
 $\forall e$  ARITH1, numbfreeocc(x1, n1, f1), x2; substr = in  $\neg$ ;
 $\forall e$  ARITH2, n1;
 $\forall e$  SUBDEFO x1, x2, f1, f2;
 $\forall e$  LEMMA1, x2, f2, n1;
 $\forall e$  LEMMA2, n1, f1, f2;
TAUTEQ  $\neg : \#2 \#1[n \leftarrow n1], \text{HPT} + 1; \neg$ ;
 $\forall i \neg, n1 \leftarrow n$ ;
TAUTEQ  $\neg : \#1, \text{HPT} + 1; \neg$ ;
 $\supset$ ! HPT,  $\neg$ ;
 $\forall i \neg, x1, x2, f1, f2$ ;

```

4.2 Printout of the proof in the many sorted logic

1 $\forall n. x. (n * (\text{len}(x) - 1)) = 0$ (1)
 2 $\forall n. (0 * n) = n$ (2)
 3 $\forall x. (\text{len}(x) - 1) = 0$ (3)
 4 $\forall n. (n - 0) = n$ (4)
 5 $\forall x. (1 \text{ gl } x) = x$ (5)
 6 **SUBT**(x,f,n) (6)
 7 **SUBT**(x,f,n) $\equiv \forall x2. k. ((k \text{ gl } x) = x2 \supset \text{FRN}(x2, n - (\text{len}(x) - k), f))$
 8 $\forall x2. k. ((k \text{ gl } x) = x2 \supset \text{FRN}(x2, n - (\text{len}(x) - k), f))$ (6)
 9 $(1 \text{ gl } x) = x \supset \text{FRN}(x, n - (\text{len}(x) - 1), f)$ (6)
 10 $(1 \text{ gl } x) = x$ (5)
 1 1 $x = x \supset \text{FRN}(x, n - (\text{len}(x) - 1), f)$ (5 6)
 1 2 $(\text{len}(x) - 1) = 0$ (3)
 1 3 $x = x \supset \text{FRN}(x, n - 0, f)$ (3 5 6)
 1 4 $(n - 0) = n$ (4)
 1 5 $x = x \supset \text{FRN}(x, n, f)$ (3 4 5 6)
 1 6 **FRN**(x,n,f) (3 4 5 6)
 1 7 **SUBT**(x,f,n) $\supset \text{FRN}(x,n,f)$ (3 4 5)
 18 $\forall x. f. n. (\text{SUBT}(x,f,n) \supset \text{FRN}(x,n,f))$ (3 4 5)
 19 **INVART**(n,f1,n,f2) $\equiv ((\text{GEB}(n \text{ gl } f2, n, f2) \equiv \text{GEB}(n \text{ gl } f1, n, f1)) \wedge ((\text{FRN}(n \text{ gl } f2, n, f2) \equiv \text{FRN}(n \text{ gl } f1, n, f1)) \wedge (n \text{ gl } f2) = (n \text{ gl } f1)))$
 20 **INVARV**(n,f1,n,f2) $\equiv ((\text{GEB}(n \text{ gl } f2, n, f2) \equiv \text{GEB}(n \text{ gl } f1, n, f1)) \wedge ((\text{FRN}(n \text{ gl } f2, n, f2) \equiv \text{FRN}(n \text{ gl } f1, n, f1)) \wedge (n \text{ gl } f2) = (n \text{ gl } f1)))$
 2 1 **INVART**(n,f1,n,f2) $\equiv \text{INVARV}(n,f1,f2)$
 22 $\forall n. f1. f2. (\text{INVART}(n,f1,n,f2) \equiv \text{INVARV}(n,f1,f2))$
 2 3 **SBT**(x1,x2,f1,f2) (2 3)
 24 **SBT**(x1,x2,f1,f2) $\equiv \forall n1. n2. (n2 = ((\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) + n1) \supset ((\neg \text{INDVAR}(n1 \text{ gl } f1) \supset (n1 \text{ gl } f1) = (n2 \text{ gl } f2)) \wedge (\text{INDVAR}(n1 \text{ gl } f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n2, f2)))))))$

25 $\forall n1. n2. (n2 = ((\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) * n1) \supset ((\neg \text{INDVAR}(n1 \text{ gl } f1) \supset$
 $(n1 \text{ gl } f1) = (n2 \text{ gl } f2)) \wedge (\text{INDVAR}(n1 \text{ gl } f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n2)) \wedge$
 $(\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n2, f2)))))) \quad (23)$

26 $n1 = ((\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) * n1) \supset ((\neg \text{INDVAR}(n1 \text{ gl } f1) \supset (n1 \text{ gl } f1) =$
 $(n1 \text{ gl } f2)) \wedge (\text{INDVAR}(n1 \text{ gl } f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n1)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset$
 $\text{INVART}(n1, f1, n1, f2)))))) \quad (23)$

27 $(\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) = 0 \quad (1)$

28 $n1 = (0 * n1) \supset ((\neg \text{INDVAR}(n1 \text{ gl } f1) \supset (n1 \text{ gl } f1) = (n1 \text{ gl } f2)) \wedge (\text{INDVAR}(n1 \text{ gl } f1) \supset$
 $((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n1)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n1, f2)))))) \quad (1 \ 23)$

29 $(0 * n1) = n1 \quad (2)$

30 $\text{SBV}(x1, x2, f1, f2) \equiv \forall n. ((\neg \text{INDVAR}(n \text{ gl } f1) \supset (n \text{ gl } f1) = (n \text{ gl } f2)) \wedge (\text{INDVAR}(n \text{ gl } f1) \supset$
 $((\text{FRN}(x1, n, f1) \supset \text{FRN}(x2, n, f2)) \wedge (\neg \text{FRN}(x1, n, f1) \supset \text{INVARV}(n, f1, f2))))))$

31 $\text{SUBT}(x2, f2, n1) \supset \text{FRN}(x2, n1, f2) \quad (3 \ 4 \ 5)$

32 $\text{INVART}(n1, f1, n1, f2) \equiv \text{INVARV}(n1, f1, f2)$

33 $(\neg \text{INDVAR}(n1 \text{ gl } f1) \supset (n1 \text{ gl } f1) = (n1 \text{ gl } f2)) \wedge (\text{INDVAR}(n1 \text{ gl } f1) \supset ((\text{FRN}(x1, n1, f1) \supset$
 $\text{FRN}(x2, n1, f2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVARV}(n1, f1, f2)))) \quad (1 \ 2 \ 3 \ 4 \ 5 \ 23)$

34 $\forall n. ((\neg \text{INDVAR}(n \text{ gl } f1) \supset (n \text{ gl } f1) = (n \text{ gl } f2)) \wedge (\text{INDVAR}(n \text{ gl } f1) \supset ((\text{FRN}(x1, n, f1) \supset$
 $\text{FRN}(x2, n, f2)) \wedge (\neg \text{FRN}(x1, n, f1) \supset \text{INVARV}(n, f1, f2)))))) \quad (1 \ 2 \ 3 \ 4 \ 5 \ 23)$

35 $\text{SBV}(x1, x2, f1, f2) \quad (1 \ 2 \ 3 \ 4 \ 5 \ 23)$

36 $\text{SBT}(x1, x2, f1, f2) \supset \text{SBV}(x1, x2, f1, f2) \quad (1 \ 2 \ 3 \ 4 \ 5)$

37 $\forall x1 \ x2 \ f1 \ f2. (\text{SBT}(x1, x2, f1, f2) \supset \text{SBV}(x1, x2, f1, f2)) \quad (1 \ 2 \ 3 \ 4 \ 5)$

4.3 FOL commands in the earlier axiomatization

```

LABEL ARITH1 ; ASSUME  $\forall n. ((\text{INTEGER}(n) \wedge \text{INDVAR}(x)) \supset (n * (\text{len}(x) - 1) = 0))$ ;
- LABEL ARITH2; ASSUME  $\forall n. (\text{INTEGER}(n) \supset (0 * n = n))$ ;
LABEL ARITH3; ASSUME  $\forall x. (\text{INDVAR}(x) \supset ((\text{len}(x) - 1) = 0))$ ;
LABEL ARITH4; ASSUME  $\forall n. (\text{INTEGER}(n) \supset (n - 0) = n)$ ;
LABEL STRING1; ASSUME  $\forall x. (\text{INDVAR}(x) \supset 1 \text{ gl } x = x)$ ;

```

Proof of the First Lemma:

$\forall x \ n \ f. ((\text{INDVAR}(x) \wedge \text{INTEGER}(n) \wedge \text{FORM}(f) \wedge \text{SUBT}(x, f, n)) \supset \text{FRN}(x, n, f))$

```

LABEL HPTLEM; ASSUME  $\text{INDVAR}(x) \wedge \text{FORM}(f) \wedge \text{INTEGER}(n) \wedge \text{SUBT}(x, f, n)$ ;
LABEL FACT ; ASSUME  $\text{INTEGER}(1)$ ;
 $\forall e \ \text{SUBSTDF } 1, x, f, n$ ;
TAUT  $-\#2\#2\#2\#2, ---, -$ ;
 $\forall e \ -, x, 1$ ;
 $\forall e \ \text{STRING1 } , x$ ; TAUT  $-\#2, \text{HPTLEM}:-;$  substr - in ---;
 $\forall e \ \text{ARITH3}, x$ ; TAUT  $-\#2, \text{HPTLEM}:-;$  substr - in ---;

```

```

Ve ARITH4,n; TAUT --: #2,HPTLEM:-;substr = in ---;
TAUTEQ FRN(x,n,f),HPTLEM:-;
>I HPTLEM,-;
LABEL LEMMA 1;Vi -,x,f,n;

```

Proof of the Second lemma : $\forall k f1 f2. (INVART(k,f1,k,f2) = INVARV(k,f1,f2))$

```

Ve SUBSTDF2,k,f1,k,f2;
Ve SUBDEF 1 ,k,f1,f2;
TAUT --: #1 = -: #1,--,--;
LABEL LEMMA2; Vi -,k,f1,f2;

```

Proof of the Main Theorem:

```

Vx1 x 2 f1 f2. ((INDVAR(x1) ^ INDVAR(x2) ^ FORM(f1) ^ FORM(f2) ^ SBT(x1,x2,f1,f2)) =>
SBV(x1,x2,f1,f2))

```

```

LABEL HPT; ASSUME INDVAR(x1) ^ INDVAR(x2) ^ FORM(f1) ^ FORM(f2) ^ SBT(x1,x2,f1,f2);

```

```

LABEL THTERM; ASSUME Vx2. (INDVAR(x2) => TERM(x2));
VE THTERM,x2;
LABEL THNFRO; ASSUME Vx1 n1 f1. INTEGER(numbfraeocc(x1,n1,f1));

```

```

Ve SUBSTDF0,x1,x2,f1,f2;
TAUT --: #2#2#2#2#2,HPT:-;
VE -,n1,n1;

```

```

LABEL AUX;ASSUME INTEGER(n1);
VE THNFRO,x1,n1,f1;

```

```

Ve ARITH1,numbfraeocc(x1,n1,f1),x2; TAUT --: #2,HPT:-;substr = in ----a
Ve ARITH2,n1;TAUT --: #2,HPT:-;SUBSTR-IN ---;
TAUTEQ --: #2,HPT:-;
Ve SUBDEFO x1,x2,f1,f2;
Ve LEMMA 1,x2,f2,n1;
Ve LEMMA2,n1,f1,f2;

```

```

- TAUTEQ ---: #2#2#1#2[n-n1], HPT :-;
>I AUX,-;
Vi -,n1;
TAUTEQ -----: #1,HPT:-;
>I HPT,-;
Vi -,x1,x2,f1,f2;

```

4.4 Printout of the proof in the earlier axiomatization

- 1 $\forall n x. ((\text{INTEGER}(n) \wedge \text{INDVAR}(x)) \Rightarrow (n * (\text{len}(x) - 1)) = 0) \quad (1)$
- 2 $\forall n. (\text{INTEGER}(n) \Rightarrow (0 * n) = n) \quad (2)$
- 3 $\forall x. (\text{INDVAR}(x) \Rightarrow (\text{len}(x) - 1) = 0) \quad (3)$

- 4 $\forall n.(\text{INTEGER}(n) \supset (n-0)=n)$ (4)
- 5 $\forall x.(\text{INDVAR}(x) \supset (1 \text{ gl } x)=x)$ (5)
- 6 $\text{INDVAR}(x) \wedge (\text{FORM}(f) \wedge (\text{INTEGER}(n) \wedge \text{SUBT}(x,f,n)))$ (6)
- 7 $\text{INTEGER}(1)$ (7)
- 8 $\text{SUBT}(x,f,n) \equiv (\text{TERM}(x) \wedge (\text{FORM}(f) \wedge (\text{INTEGER}(n) \wedge \forall x2.k.((\text{INDVAR}(x2) \wedge (\text{INTEGER}(k) \wedge (k \text{ gl } x) = x2)) \supset \text{FRN}(x2,n-(\text{len}(x)-k),f))))))$
- 9 $\forall x2.k.((\text{INDVAR}(x2) \wedge (\text{INTEGER}(k) \wedge (k \text{ gl } x)=x2)) \supset \text{FRN}(x2,n-(\text{len}(x)-k),f))$ (6)
- 10 $(\text{INDVAR}(x) \wedge (\text{INTEGER}(1) \wedge (1 \text{ gl } x)=x)) \supset \text{FRN}(x,n-(\text{len}(x)-1),f)$ (6)
- 11 $\text{INDVAR}(x) \supset (1 \text{ gl } x)=x$ (5 9)
- 12 $(1 \text{ gl } x)=x$ (5 6 7)
- 13 $(\text{INDVAR}(x) \wedge (\text{INTEGER}(1) \wedge x=x)) \supset \text{FRN}(x,n-(\text{len}(x)-1),f)$ (5 6 7)
- 14 $\text{INDVAR}(x) \supset (\text{len}(x)-1)=0$ (3)
- 15 $(\text{len}(x)-1)=0$ (3 5 6 7)
- 16 $(\text{INDVAR}(x) \wedge (\text{INTEGER}(1) \wedge x=x)) \supset \text{FRN}(x,n-0,f)$ (3 5 6 7)
- 17 $\text{INTEGER}(n) \supset (n-0)=n$ (4)
- 18 $(n-0)=n$ (3 4 5 6 7)
- 19 $(\text{INDVAR}(x) \wedge (\text{INTEGER}(1) \wedge x=x)) \supset \text{FRN}(x,n,f)$ (3 4 5 6 7)
- 20 $\text{FRN}(x,n,f)$ (3 4 5 6 7)
- 21 $(\text{INDVAR}(x) \wedge (\text{FORM}(f) \wedge (\text{INTEGER}(n) \wedge \text{SUBT}(x,f,n)))) \supset \text{FRN}(x,n,f)$ (3 4 5 7)
- 22 $\forall x.f.n.((\text{INDVAR}(x) \wedge (\text{FORM}(f) \wedge (\text{INTEGER}(n) \wedge \text{SUBT}(x,f,n)))) \supset \text{FRN}(x,n,f))$ (3 4 5 7)
- 23 $\text{INVART}(k,f1,k,f2) \equiv (\text{INTEGER}(k) \wedge (\text{FORM}(f1) \wedge (\text{INTEGER}(k) \wedge (\text{FORM}(f2) \wedge ((\text{GEB}(k \text{ gl } f2,k,f2) \equiv \text{GEB}(k \text{ gl } f1,k,f1)) \wedge ((\text{FRN}(k \text{ gl } f2,k,f2) \equiv \text{FRN}(k \text{ gl } f1,k,f1)) \wedge (k \text{ gl } f2)=(k \text{ gl } f1)))))))$
- 24 $\text{INVARV}(k,f1,f2) \equiv (\text{INTEGER}(k) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(f2) \wedge ((\text{GEB}(k \text{ gl } f2,k,f2) \equiv \text{GEB}(k \text{ gl } f1,k,f1)) \wedge ((\text{FRN}(k \text{ gl } f2,k,f2) \equiv \text{FRN}(k \text{ gl } f1,k,f1)) \wedge (k \text{ gl } f2)=(k \text{ gl } f1))))))$
- 25 $\text{INVART}(k,f1,k,f2) \equiv \text{INVARV}(k,f1,f2)$
- 26 $\forall k.f1.f2.(\text{INVART}(k,f1,k,f2) \equiv \text{INVARV}(k,f1,f2))$
- 27 $\text{INDVAR}(x1) \wedge (\text{INDVAR}(x2) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(f2) \wedge \text{SBT}(x1,x2,f1,f2))))$ (27)
- 28 $\forall x2.(\text{INDVAR}(x2) \supset \text{TERM}(x2))$ (28)

29 $\text{INDVAR}(x2) \supset \text{TERM}(x2)$ (28)

30 $\forall x1 \, n1 \, f1. \text{INTEGER}(\text{numbfreeocc}(x1, n1, f1))$ (30)

31 $\text{SBT}(x1, x2, f1, f2) \equiv ((\text{INDVAR}(x1) \wedge (\text{TERM}(x2) \wedge (\text{FORM}(f1) \wedge \text{FORM}(f2)))) \supset \forall n1 \, n2. ((\text{INTEGER}(n1) \wedge (\text{INTEGER}(n2) \wedge n2 = ((\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) + n1))) \supset ((\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n2 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n2, f2)))))))$

32 $\forall n1 \, n2. ((\text{INTEGER}(n1) \wedge (\text{INTEGER}(n2) \wedge n2 = ((\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) + n1))) \supset ((\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n2 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n2, f2))))))$ (27 28 30)

33 $(\text{INTEGER}(n) \wedge (\text{INTEGER}(n1) \wedge n1 = ((\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) + n1))) \supset ((\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n1 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n1)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n1, f2))))$ (27 28 30)

34 $\text{INTEGER}(n1)$ (34)

35 $\text{INTEGER}(\text{numbfreeocc}(x1, n1, f1))$ (30)

36 $(\text{INTEGER}(\text{numbfreeocc}(x1, n1, f1)) \wedge \text{INDVAR}(x2)) \supset (\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) = 0$ (1)

37 $(\text{numbfreeocc}(x1, n1, f1) * (\text{len}(x2) - 1)) = 0$ (1 27 28 30 34)

38 $(\text{INTEGER}(n1) \wedge (\text{INTEGER}(n1) \wedge n1 = (0 * n1))) \supset ((\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n1 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n1)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n1, f2))))$ 0 27 28 30 34)

39 $\text{INTEGER}(n1) \supset (0 * n1) = n1$ (2)

40 $(0 * n1) = n1$ (1 2 27 28 30 34)

41 $(\text{INTEGER}(n1) \wedge (\text{INTEGER}(n1) \wedge n1 = n1)) \supset ((\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n1 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n1)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n1, f2))))$ (1 2 27 28 30 34)

42 $(\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n1 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{SUBT}(x2, f2, n1)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVART}(n1, f1, n1, f2))))$ (1 2 27 28 30 34)

43 $\text{SBV}(x1, x2, f1, f2) \equiv ((\text{INDVAR}(x1) \wedge (\text{INDVAR}(x2) \wedge (\text{FORM}(f1) \wedge \text{FORM}(f2)))) \supset \forall n. (\text{INTEGER}(n) \supset ((\neg \text{INDVAR}(n \, gl \, f1) \supset (n \, gl \, f1) = (n \, gl \, f2)) \wedge (\text{INDVAR}(n \, gl \, f1) \supset ((\text{FRN}(x1, n, f1) \supset \text{FRN}(x2, n, f2)) \wedge (\neg \text{FRN}(x1, n, f1) \supset \text{INVARY}(n, f1, f2)))))))$

44 $(\text{INDVAR}(x2) \wedge (\text{FORM}(f2) \wedge (\text{INTEGER}(n1) \wedge \text{SUBT}(x2, f2, n1)))) \supset \text{FRN}(x2, n1, f2)$ (3 4 5 7)

45 $\text{INVART}(n1, f1, n1, f2) \equiv \text{INVARY}(n1, f1, f2)$

46 $(\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n1 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{FRN}(x2, n1, f2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVARY}(n1, f1, f2))))$ (1 2 3 4 5 7 27 28 30 34)

47 $\text{INTEGER}(n1) \supset ((\neg \text{INDVAR}(n1 \, gl \, f1) \supset (n1 \, gl \, f1) = (n1 \, gl \, f2)) \wedge (\text{INDVAR}(n1 \, gl \, f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{FRN}(x2, n1, f2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVARY}(n1, f1, f2))))$ (1 2 3 4 5

- 7 27 28 30)
- 48 $\forall n1. (\text{INTEGER}(n1) \supset ((\neg \text{INDVAR}(n1 \text{ gl } f1) \supset (n1 \text{ gl } f1) = (n1 \text{ gl } f2)) \wedge (\text{INDVAR}(n1 \text{ gl } f1) \supset ((\text{FRN}(x1, n1, f1) \supset \text{FRN}(x2, n1, f2)) \wedge (\neg \text{FRN}(x1, n1, f1) \supset \text{INVARV}(n1, f1, f2)))))) \quad (1 \ 2 \ 3 \ 4 \ 5 \ 7 \ 27 \ 28 \ 30)$
- 49 **SBV**(x1,x2,f1,f2) (1 2 3 4 5 7 27 28 30 34)
- 50 $(\text{INDVAR}(x1) \wedge (\text{INDVAR}(x2) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(f2) \wedge \text{SBT}(x1, x2, f1, f2)))) \supset \text{SBV}(x1, x2, f1, f2)) \quad (1 \ 2 \ 3 \ 4 \ 5 \ 7 \ 28 \ 30 \ 34)$
- 51 $\forall x1 \ x2 \ f1 \ f2. ((\text{INDVAR}(x1) \wedge (\text{INDVAR}(x2) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(f2) \wedge \text{SBT}(x1, x2, f1, f2)))) \supset \text{SBV}(x1, x2, f1, f2)) \quad (1 \ 2 \ 3 \ 4 \ 5 \ 7 \ 28 \ 30)$

APPENDIX 5

THE PROOF THAT UNIVERSAL QUANTIFIER CAN BE INTERCHANGED

5.1 FOL commands for the main lemma in the many sorted logic

```

LABEL TH1 ; ASSUME  $\forall x_1 x_2 f_1 f_2. (SBT(x_1, x_2, f_1, f_2) \supset SBV(x_1, x_2, f_1, f_2))$ ;
 $\forall e$  TH1,  $x, x, f_1, sbt(x, x, f_1)$ ;
VE SUBSTDF3  $x, x, f_1, sbt(x, x, f_1)$ ;
 $\forall e$  SUBDEFO  $x, x, f_1, sbt(x, x, f_1)$ ;
tauteq  $\neg : \#2, 1 :-$ ;
 $\forall e$   $\neg, n$ ;
VE FREEVO,  $x, n, f_1$ ;
VE FREEVO,  $x, n, sbt(x, x, f_1)$ ;
VE SUBDEF 1  $n, f_1, sbt(x, x, f_1)$ ;
tauteq  $(n \text{ gl } f_1) = (n \text{ gl } sbt(x, x, f_1)), 11, 17, 18$ ;
Vi  $\neg, n$ ;
VE EQS  $f_1, sbt(x, x, f_1)$ ;
tauteq  $sbt(x, x, f_1) = f_1, --, -$ ;
Vi  $\neg, x, f_1 \leftarrow f$ ;

```

5.2 Printout of the proof in the many sorted logic

- 1 $\forall x_1 x_2 f_1 f_2. (SBT(x_1, x_2, f_1, f_2) \supset SBV(x_1, x_2, f_1, f_2)) (1)$
- 2 $SBT(x, x, f_1, sbt(x, x, f_1)) \supset SBV(x, x, f_1, sbt(x, x, f_1)) (1)$
- 3 $SBT(x, x, f_1, sbt(x, x, f_1)) \neg sbt(x, x, f_1) = sbt(x, x, f_1)$
- 4 $SBV(x, x, f_1, sbt(x, x, f_1)) = \forall n. ((\neg INDVAR(n \text{ gl } f_1) \supset (n \text{ gl } f_1) = (n \text{ gl } sbt(x, x, f_1))) \wedge (INDVAR(n \text{ gl } f_1) \supset ((FRN(x, n, f_1) \supset FRN(x, n, sbt(x, x, f_1))) \wedge (\neg FRN(x, n, f_1) \supset INVARV(n, f_1, sbt(x, x, f_1))))))$
- 5 $\forall n. ((\neg INDVAR(n \text{ gl } f_1) \supset (n \text{ gl } f_1) = (n \text{ gl } sbt(x, x, f_1))) \wedge (INDVAR(n \text{ gl } f_1) \supset ((FRN(x, n, f_1) \supset FRN(x, n, sbt(x, x, f_1))) \wedge (\neg FRN(x, n, f_1) \supset INVARV(n, f_1, sbt(x, x, f_1)))))) (1)$
- 6 $(\neg INDVAR(n \text{ gl } f_1) \supset (n \text{ gl } f_1) = (n \text{ gl } sbt(x, x, f_1))) \wedge (INDVAR(n \text{ gl } f_1) \supset ((FRN(x, n, f_1) \supset FRN(x, n, sbt(x, x, f_1))) \wedge (\neg FRN(x, n, f_1) \supset INVARV(n, f_1, sbt(x, x, f_1)))))) (1)$
- 7 $FRN(x, n, f_1) = (x = (n \text{ gl } f_1) \wedge \neg GEB(x, n, f_1)) \quad \text{VE FREEVO } x, n, f_1$
- 8 $FRN(x, n, sbt(x, x, f_1)) = (x = (n \text{ gl } sbt(x, x, f_1)) \wedge \neg GEB(x, n, sbt(x, x, f_1)))$
- 9 $INVARV(n, f_1, sbt(x, x, f_1)) = ((GEB(n \text{ gl } sbt(x, x, f_1), n, sbt(x, x, f_1)) = GEB(n \text{ gl } f_1, n, f_1)) \wedge ((FRN(n \text{ gl } sbt(x, x, f_1), n, sbt(x, x, f_1)) = FRN(n \text{ gl } f_1, n, f_1)) \wedge (n \text{ gl } sbt(x, x, f_1) = (n \text{ gl } f_1))))$
- 10 $(n \text{ gl } f_1) = (n \text{ gl } sbt(x, x, f_1)) (1)$


```

∀e DEPEND0 f cc sq,sq,f 1;
UNIFY --:#2#2#2#2#2,GENE1;

TAUTEQ DEPEND(f cc sq,f 1)  $\supset$  AXIOM (f 1),1:-; .
Vi -,f1 $\leftarrow$ f1;
TAUTEQ THN1:#2 = THN1:#1,THN1;
Λi PROOFTR, -, -;
LABEL USEFUL;
∀e PROVABLE f;
UNIFY --:#2,-;
TAUT --:#1,1:-;
LABEL CI TH1;
⊃i HPT,-;
Proof of the Lemma: BEW(f)  $\supset$  BEW(x gen f)

LABEL HPT1;
ASSUME BEW(f);

TAUT USEFUL:#2, -,HPT 1,USEFUL;
∃e -,sq;

Λe --:#2#2;
∀e -, f1;
∀e GENRUL2 x,sq;
∀e THEORY x,f1;
TAUTEQ --:#2#1#1[f $\leftarrow$ f1],HPT1:-;
Vi -,f1 $\leftarrow$ f1;
TAUT ----:#1,HPT1:-;

∀e GENRUL1((x gen f) cc sq) , sq ,x,x ;
LABEL THN2;
∀e THEON1 x gen f , sq ;
∀e THEON2 x gen f , sq ;
TAUTEQ ----:#2#2#2#1[f1  $\leftarrow$  f] , THTAUT,HPT1:-;
UNIFY ----:#2#2#2 , -;
TAUTEQ -----:#1 , THTAUT,HPT1:-;

∀e PROOF (x gen f) cc sq ;
LABEL GEN1;
vi -- , GENE((x gen f) cc sq,sq,x,x) , EXI((x gen f) cc sq,sq,x,x) ;
UNIFY --:#2#2#2#1 , - ;

LABEL PROOFTR 1;
TAUT ----:#1,HPT1:-,THTAUT;

∀e DEPEND0 (x gen f) cc sq, sq,f 1;
31 GEN1 ,x $\leftarrow$ t OCC 3 6 9,x $\leftarrow$ x1 OCC 2 4 6;

TAUTEQ DEPEND((x gen f) cc sq,f 1)  $\supset$  AXIOM (f 1) ,THTAUT,HPT1:-;

```

```

Vi -,f1←f1;
TAUT EQ THN2:#2 = THN2:#1,THN2;
Ai PROOFTRI , -, -- ;
V● PROVABLE x gen f;
UNIFY -:#2,--;
TAUT ---:# 1 ,THTAUT,HPT1:-;
LABEL C2TH1;
⇒I HPT 1,-;
⇒I C1TH1,C2TH1;
LABEL TH1;
Vi -,x,f;
V● TH1 x1,x2 gen f;
V e TH1 x2,f;
V● TH1 x1,f;
V● TH1 x2,x1 gen f;
TAUT ----:# 1 ⇒-:# 1, TH1:-;
vi -,x 1 ,x2,f;

```

5.4 Printout of the proof of the theorem in the marry sorted logic

```

1  V x f.sbt(x,x,f)=f (1)
2  V f sq.scar(f cc sq)=f (2)
3  V f sq.scdr(f cc sq)=sq (3)
4  BEW(x gen f) (4)
5  sbt(x,x,f)=f (1)
6  BEW(x gen f)⇒∃sq.(PROOFTREE(sq)∧((x gen f)=scar(sq)∧∀f 1 .(DEPEND(sq,f 1)⇒AXIOM(f 1))))
7  ∃sq.(PROOFTREE(sq)∧((x gen f)=scar(sq)∧∀f 1 .(DEPEND(sq,f 1)⇒AXIOM(f 1)))) (4)
8  PROOFTREE(sq)∧((x gen f)=scar(sq)∧∀f 1 .(DEPEND(sq,f 1)⇒AXIOM(f 1))) (8)
9  GENE(f cc sq,sq,x,x)⇒(scdr(f cc sq)=sq∧(PROOFTREE(sq)∧∃f 1 .(scar(sq)=(x gen f 1)∧
  scar(f cc sq)=sbt(x,x,f 1))))
10 scar(f cc sq)=f (2)
11 scdr(f cc sq)=sq (3)
12 scar(sq)=(x gen f)∧scar(f cc sq)=sbt(x,x,f) (1 2 3 4 8)
13 ∃f 1 .(scar(sq)=(x gen f 1)∧scar(f cc sq)=sbt(x,x,f 1)) (1 2 3 4 8)
14 GENE(f cc sq,sq,x,x) (1 2 3 4 8)
15 PROOFTREE(f cc sq)⇒(FORM(f cc sq)∨(∃pf.(ORI(f cc sq,pf)∨(ANDE(f cc sq,pf)∨
  (FALSEE(f cc sq,pf)∨(NOTI(f cc sq,pf)∨(NOTE(f cc sq,pf)∨IMPLI(f cc sq,pf)))))))))v

```

$(3pf \ x \ t.(GENI(f \ cc \ sq, pf, x, t) \vee (GENE(f \ cc \ sq, pf, x, t) \vee EXI(f \ cc \ sq, pf, x, t))) \vee$
 $(3pf \ 1 \ pf2.(ANDI(f \ cc \ sq, pf1, pf2) \vee (FALSEI(f \ cc \ sq, pf1, pf2) \vee IMPLI(f \ cc \ sq, pf1, pf2))) \vee$
 $3pf1 \ pf2 \ x \ t.EXE(f \ cc \ sq, pf1, pf2, x, t) \vee 3pf1 \ pf2 \ pf3.ORE(f \ cc \ sq, pf1, pf2, pf3))))))$

16 $GENI(f \ cc \ sq, sq, x, x) \vee (GENE(f \ cc \ sq, sq, x, x) \vee EXI(f \ cc \ sq, sq, x, x)) \quad (1 \ 2 \ 3 \ 4 \ 8)$

17 $3pf \ x \ t.(GENI(f \ cc \ sq, pf, x, t) \vee (GENE(f \ cc \ sq, pf, x, t) \vee EXI(f \ cc \ sq, pf, x, t))) \quad (1 \ 2 \ 3 \ 4 \ 8)$

18 $PROOFTREE(f \ cc \ sq) \quad (1 \ 2 \ 3 \ 4 \ 8)$

19 $\forall f1.(DEPEND(sq, f1) \supset AXIOM(f1)) \quad (8)$

20 $DEPEND(sq, f1) \supset AXIOM(f1) \quad (8)$

21 $PROOFTREE(f \ cc \ sq) \supset (PROOFTREE(sq) \supset ((sq = scdr(f \ cc \ sq) \supset (DEPEND(f \ cc \ sq, f1) \supset$
 $DEPEND(sq, f1))) \supset (ORI(f \ cc \ sq, sq) \vee (ANDE(f \ cc \ sq, sq) \vee (FALSEE(f \ cc \ sq, sq) \vee$
 $(3f1.(NOTID(f \ cc \ sq, sq, f1) \vee (NOTED(f \ cc \ sq, sq, f1) \vee IMPLID(f \ cc \ sq, sq, f1))) \wedge f1 \neq f1) \vee$
 $3x \ t.(GENI(f \ cc \ sq, sq, x, t) \vee (GENE(f \ cc \ sq, sq, x, t) \vee EXI(f \ cc \ sq, sq, x, t)))))))$

22 $\exists x \ t.(GENI(f \ cc \ sq, sq, x, t) \vee (GENE(f \ cc \ sq, sq, x, t) \vee EXI(f \ cc \ sq, sq, x, t))) \quad (1 \ 2 \ 3 \ 4 \ 8)$

23 $DEPEND(f \ cc \ sq, f1) \supset AXIOM(f1) \quad (1 \ 2 \ 3 \ 4 \ 8)$

24 $\forall f1.(DEPEND(f \ cc \ sq, f1) \supset AXIOM(f1)) \quad (1 \ 2 \ 3 \ 4 \ 8)$

25 $f = scar(f \ cc \ sq) \quad (2)$

26 $PROOFTREE(f \ cc \ sq) \wedge (f = scar(f \ cc \ sq) \wedge \forall f1.(DEPEND(f \ cc \ sq, f1) \supset AXIOM(f1))) \quad (1 \ 2 \ 3 \ 4 \ 8)$

27 $BEW(f) \equiv \exists sq.(PROOFTREE(sq) \wedge (f = scar(sq) \wedge \forall f1.(DEPEND(sq, f1) \supset AXIOM(f1))))$

28 $\exists sq.(PROOFTREE(sq) \wedge (f = scar(sq) \wedge \forall f1.(DEPEND(sq, f1) \supset AXIOM(f1)))) \quad (1 \ 2 \ 3 \ 4)$

29 $BEW(f) \quad (1 \ 2 \ 3 \ 4)$

30 $BEW(x \ gen \ f) \supset BEW(f) \quad (1 \ 2 \ 3)$

31 $BEW(f) \quad (3 \ 1)$

32 $\exists sq.(PROOFTREE(sq) \wedge (f = scar(sq) \wedge \forall f1.(DEPEND(sq, f1) \supset AXIOM(f1)))) \quad (31)$

33 $PROOFTREE(sq) \wedge (f = scar(sq) \wedge \forall f1.(DEPEND(sq, f1) \supset AXIOM(f1))) \quad (33)$

34 $\forall f1.(DEPEND(sq, f1) \supset AXIOM(f1)) \quad (33)$

35 $DEPEND(sq, f1) \supset AXIOM(f1) \quad (33)$

36 $APGENI(x, sq) \equiv (\forall f1.(DEPEND(sq, f1) \supset \neg FR(x, f1)) \wedge PROOFTREE(sq))$

37 $AXIOM(f1) \supset \neg FR(x, f1)$

38 $DEPEND(sq, f1) \supset \neg FR(x, f1) \quad (3 \ 1 \ 33)$

```

39  $\forall f1. (DEPEND(sq, f1) \supset \neg FR(x, f1))$  (3 1 33)

40 APGENI(x, sq) (3 1 33)

41 GENI((x gen f) cc sq, sq, x, x) = (scdr((x gen f) cc sq) = sq  $\wedge$  (PROOFTREE(sq)  $\wedge$ 
  3f1. (scar((x gen f) cc sq) = (x gen f 1)  $\wedge$  (scar(sq) = sbt(x, x, f1)  $\wedge$  APGENI(x, sq))))))

42 scar((x gen f) cc sq) = (x gen f) (2)

43 scdr((x gen f) cc sq) = sq (3)

44 scar((x gen f) cc sq) = (x gen f)  $\wedge$  (scar(sq) = sbt(x, x, f)  $\wedge$  APGENI(x, sq)) (1 2 3 31 33)

45 3f1. (scar((x gen f) cc sq) = (x gen f 1)  $\wedge$  (scar(sq) = sbt(x, x, f1)  $\wedge$  APGENI(x, sq))) (1 2 3 31 33)

46 GENI((x gen f) cc sq, sq, x, x) (1 2 3 31 33)

47 PROOFTREE((x gen f) cc sq) = (FORM((x gen f) cc sq)  $\vee$  (3pf. (ORI((x gen f) cc sq, pf)  $\vee$ 
  (ANDE((x gen f) cc sq, pf)  $\vee$  (FALSEE((x gen f) cc sq, pf)  $\vee$  (NOTI((x gen f) cc sq, pf)  $\vee$ 
  (NOTE((x gen f) cc sq, pf)  $\vee$  IMPLI((x gen f) cc sq, pf))))))  $\vee$  (3pf x1 t. (GENI((x gen f)
  cc sq, pf, x1, t)  $\vee$  (GENE((x gen f) cc sq, pf, x1, t)  $\vee$  EXI((x gen f) cc sq, pf, x1, t)))  $\vee$ 
  (3pf1 pf2. (ANDI((x gen f) cc sq, pf1, pf2)  $\vee$  (FALSEI((x gen f) cc sq, pf1, pf2)  $\vee$ 
  IMPLI((x gen f) cc sq, pf1, pf2)))  $\vee$  (3pf1 pf2 x1 t. EXE((x gen f) cc sq, pf1, pf2, x1, t)  $\vee$ 
  3pf1 pf2 pf3. ORE((x gen f) cc sq, pf1, pf2, pf3))))))

48 GENI((x gen f) cc sq, sq, x, x)  $\vee$  (GENE((x gen f) cc sq, sq, x, x)  $\vee$  EXI((x gen f) cc sq,
  sq, x, x)) (1 2 3 31 33)

49 3pf x1 t. (GENI((x gen f) cc sq, pf, x1, t)  $\vee$  (GENE((x gen f) cc sq, pf, x1, t)  $\vee$ 
  EXI((x gen f) cc sq, pf, x1, t))) (1 2 3 31 33)

50 PROOFTREE((x gen f) cc sq) (1 2 3 31 33)

51 PROOFTREE((x gen f) cc sq) = (PROOFTREE(sq) = ((sq = scdr((x gen f) cc sq) = (DEPEND(
  (x gen f) cc sq, f1) = DEPEND(sq, f1))) = (ORI((x gen f) cc sq, sq)  $\vee$  (ANDE((x gen f)
  cc sq, sq)  $\vee$  (FALSEE((x gen f) cc sq, sq)  $\vee$  (3f. ((NOTID((x gen f) cc sq, sq, f)  $\vee$ 
  (NOTED((x gen f) cc sq, sq, f)  $\vee$  IMPLID((x gen f) cc sq, sq, f)))  $\wedge$  f1)  $\vee$ 
  3x1 t. (GENI((x gen f) cc sq, sq, x1, t)  $\vee$  (GENE((x gen f) cc sq, sq, x1, t)  $\vee$  EXI(
  (x gen f) cc sq, sq, x1, t))))))))))

52 3x1 t. (GENI((x gen f) cc sq, sq, x1, t)  $\vee$  (GENE((x gen f) cc sq, sq, x1, t)  $\vee$ 
  EXI((x gen f) cc sq, sq, x1, t))) (1 2 3 31 33)

53 DEPEND((x gen f) cc sq, f1) = AXIOM(f1) (1 2 3 31 33)

54  $\forall f1. (DEPEND((x gen f) cc sq, f1) = \text{AXIOM}(f1))$  (1 2 3 31 33)

55 (x gen f) = scar((x gen f) cc sq) (2)

56 PROOFTREE((x gen f) cc sq)  $\wedge$  ((x gen f) = scar((x gen f) cc sq)  $\wedge$ 
   $\forall f1. (DEPEND((x gen f) cc sq, f1) = \text{AXIOM}(f1)))$  (1 2 3 31 33)

57 BEW(x gen f) = 3sq. (PROOFTREE(sq)  $\wedge$  ((x gen f) = scar(sq)  $\wedge$   $\forall f1. (DEPEND(sq, f1) = \text{AXIOM}(f1))))$ 

```

```

58 3sq.(PROOFTREE(sq)^(x 0)ascar(sq)^(Vf 1 .(DEPEND(sq,f 1)=AXIOM(f 1)))) (1 2 3 31)
59 BEW(x gen f) (1 2 3 31)
60 BEW(f)⊃BEW(x gen f) (1 2 3)
61 BEW(x 0)⊃BEW(f) (1 2 3)
62 Vx f.(BEW(x gen f)⊃BEW(f)) (1 2 3)
63 BEW(x1 gen (x2 gen f))⊃BEW(x2 gen f) (1 2 3)
64 BEW(x2 0)⊃BEW(f) (1 2 3)
65 BEW(x1 f)⊃BEW(f) (1 2 3)
66 BEW(x2 gen (x1 gen f))⊃BEW(x1 gen f) (1 2 3)
67 BEW(x1 gen (x2 gen f))⊃BEW(x2 gen (x1 gen f)) (1 2 3)
68 Vx 1 x2 f.(BEW(x1 gen (x2 gen f))⊃BEW(x2 gen (x1 gen f))) (1 2 3)

```

5.5 FOL commands for the main lemma in the earlier axiomatization

```

LABEL HPT; ASSUME INDVAR(x) A FORM(f1);
LABEL TH1; ASSUME Vx1 x2 f 1 f2.((INDVAR(x1) ^ INDVAR(x2) ^ FORM(f 1) A FORM(f2) A
    SBT(x1,x2,f1,f2))⊃SBV(x1,x2,f 1 ,f2));
LABEL TH2; ASSUME Vx.(INDVAR(x)⊃TERM(x));
LABEL TH3; ASSUME Vx.(FORM(x)⊃STRING(x));
V0 TH1, x,x,f 1 ,sbt(x,x,f 1);
V0 TH2, x;
V0 TH3, f1;
V0 TH3, sbt(x,x,f 1);
VE SUBSTDF3 x,x,f 1 ,sbt(x,x,f 1);
VE SUBSTDF4 x,x, f 1;
V0 SUBDEFO x, x,f 1 ,sbt(x,x,f 1);
tauteq -:#2#2,l:-;
V0 -,n;
VE FREEVO, x, n, f 1;
VE FREEVO, x, n, sbt(x,x,f 1);
VE SUBDEF1 n,f 1 ,sbt(x,x,f 1);
tauteq INTEGER(n) ⊃ ((n gl f1)=(n gl sbt(x,x,f 1))) l:-;
Vi -,n;
VE EQS, f 1,sbt(x,x,f 1);
taut -:#2#2,l:-;
⊃i,l:-;
Vl -,x,f 1←f;

```

5.6 Printout of the proof of the main lemma in the second axiomatization

- 1 $\text{INDVAR}(x) \wedge \text{FORM}(f1) \quad (1) \text{ ASSUME}$
- 2 $\forall x1 \ x2 \ f1 \ f2. ((\text{INDVAR}(x1) \wedge (\text{INDVAR}(x2) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(f2) \wedge \text{SBT}(x1, x2, f1, f2)))))) \Rightarrow \text{SBV}(x1, x2, f1, f2)) \quad (2) \text{ ASSUME}$
- 3 $\forall x. (\text{INDVAR}(x) \Rightarrow \text{TERM}(x)) \quad (3) \text{ ASSUME}$
- 4 $\forall x. (\text{FORM}(x) \Rightarrow \text{STRING}(x)) \quad (4) \text{ ASSUME}$
- 5 $(\text{INDVAR}(x) \wedge (\text{INDVAR}(x) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(\text{sbt}(x, x, f1)) \wedge \text{SBT}(x, x, f1, \text{sbt}(x, x, f1)))))) \Rightarrow \text{SBV}(x, x, f1, \text{sbt}(x, x, f1)) \quad (2) \forall E \ 2 \ x, x, f1, \text{sbt}(x, x, f1)$
- 6 $\text{INDVAR}(x) \Rightarrow \text{TERM}(x) \quad (3) \forall E \ 3 \ x$
- 7 $\text{FORM}(f1) \Rightarrow \text{STRING}(f1) \quad (4) \forall E \ 4 \ f1$
- 8 $\text{FORM}(\text{sbt}(x, x, f1)) \Rightarrow \text{STRING}(\text{sbt}(x, x, f1)) \quad (4) \forall E \ 4 \ \text{sbt}(x, x, f1)$
- 9 $(\text{INDVAR}(x) \wedge (\text{TERM}(x) \wedge (\text{FORM}(f1) \wedge \text{FORM}(\text{sbt}(x, x, f1)))) \Rightarrow (\text{SBT}(x, x, f1, \text{sbt}(x, x, f1)) \Rightarrow \text{sbt}(x, x, f1) = \text{sbt}(x, x, f1)) \quad \text{VE SUBSTDF3 } x, x, f1, \text{sbt}(x, x, f1)$
- 10 $(\text{INDVAR}(x) \wedge (\text{TERM}(x) \wedge \text{FORM}(f1))) \Rightarrow \text{FORM}(\text{sbt}(x, x, f1)) \quad \text{VE SUBSTDF4 } x, x, f1$
- 11 $\text{SBV}(x, x, f1, \text{sbt}(x, x, f1)) \Rightarrow ((\text{INDVAR}(x) \wedge (\text{INDVAR}(x) \wedge (\text{FORM}(f1) \wedge \text{FORM}(\text{sbt}(x, x, f1)))) \Rightarrow \forall n. (\text{INTEGER}(n) \Rightarrow ((\neg \text{INDVAR}(n \text{ gl } f1) \Rightarrow (n \text{ gl } f1) = (n \text{ gl } \text{sbt}(x, x, f1))) \wedge (\text{INDVAR}(n \text{ gl } f1) \Rightarrow ((\text{FRN}(x, n, f1) \Rightarrow \text{FRN}(x, n, \text{sbt}(x, x, f1))) \wedge (\neg \text{FRN}(x, n, f1) \Rightarrow \text{INVARV}(n, f1, \text{sbt}(x, x, f1))))))) \Rightarrow \text{INVARV}(n, f1, \text{sbt}(x, x, f1)))) \quad (1 \ 2 \ 3 \ 4) \ 1 : 11$
- 12 $\forall n. (\text{INTEGER}(n) \Rightarrow ((\neg \text{INDVAR}(n \text{ gl } f1) \Rightarrow (n \text{ gl } f1) = (n \text{ gl } \text{sbt}(x, x, f1))) \wedge (\text{INDVAR}(n \text{ gl } f1) \Rightarrow ((\text{FRN}(x, n, f1) \Rightarrow \text{FRN}(x, n, \text{sbt}(x, x, f1))) \wedge (\neg \text{FRN}(x, n, f1) \Rightarrow \text{INVARV}(n, f1, \text{sbt}(x, x, f1))))))) \quad (1 \ 2 \ 3 \ 4) \ 1 : 11$
- 13 $\text{INTEGER}(n) \Rightarrow ((\neg \text{INDVAR}(n \text{ gl } f1) \Rightarrow (n \text{ gl } f1) = (n \text{ gl } \text{sbt}(x, x, f1))) \wedge (\text{INDVAR}(n \text{ gl } f1) \Rightarrow ((\text{FRN}(x, n, f1) \Rightarrow \text{FRN}(x, n, \text{sbt}(x, x, f1))) \wedge (\neg \text{FRN}(x, n, f1) \Rightarrow \text{INVARV}(n, f1, \text{sbt}(x, x, f1))))))) \quad (1 \ 2 \ 3 \ 4) \ \text{VE } 12 \ n$
- 14 $\text{FRN}(x, n, f1) \Rightarrow (x = (n \text{ gl } f1) \wedge \neg \text{GEB}(x, n, f1)) \quad \text{VE FREEVO } x, n, f1$
- 15 $\text{FRN}(x, n, \text{sbt}(x, x, f1)) \Rightarrow (x = (n \text{ gl } \text{sbt}(x, x, f1)) \wedge \neg \text{GEB}(x, n, \text{sbt}(x, x, f1))) \quad \text{VE FREEVO } x, n, \text{sbt}(x, x, f1)$
- 16 $\text{INVARV}(n, f1, \text{sbt}(x, x, f1)) \Rightarrow (\text{INTEGER}(n) \wedge (\text{FORM}(f1) \wedge (\text{FORM}(\text{sbt}(x, x, f1)) \wedge ((\text{GEB}(n \text{ gl } \text{sbt}(x, x, f1), n, \text{sbt}(x, x, f1)) \Rightarrow \text{GEB}(n \text{ gl } f1, n, f1)) \wedge ((\text{FRN}(n \text{ gl } \text{sbt}(x, x, f1), n, \text{sbt}(x, x, f1)) \Rightarrow \text{FRN}(n \text{ gl } f1, n, f1)) \wedge (n \text{ gl } \text{sbt}(x, x, f1) = (n \text{ gl } f1)))))) \quad \text{VE SUBDEF1 } n, f1, \text{sbt}(x, x, f1)$
- 17 $\text{INTEGER}(n) \Rightarrow (n \text{ gl } f1) = (n \text{ gl } \text{sbt}(x, x, f1)) \quad (1 \ 2 \ 3 \ 4) \ 1 : 16$
- 18 $\forall n. (\text{INTEGER}(n) \Rightarrow (n \text{ gl } f1) = (n \text{ gl } \text{sbt}(x, x, f1))) \quad (1 \ 2 \ 3 \ 4) \ \text{VI } 17 \ n \leftarrow n$
- 19 $(\text{STRING}(f1) \wedge \text{STRING}(\text{sbt}(x, x, f1))) \Rightarrow (\forall n. (\text{INTEGER}(n) \Rightarrow (n \text{ gl } f1) = (n \text{ gl } \text{sbt}(x, x, f1))) \Rightarrow f1 = \text{sbt}(x, x, f1)) \quad \text{VE EQS } f1, \text{sbt}(x, x, f1)$

5.7 FOL commands in the earlier axiomatization

Proof of the Lemma $\mathbf{BEW}(x \text{ gen } f) \supset \mathbf{BEW}(f)$ Under the Assumption: $\mathbf{INDVAR}(x) \wedge \mathbf{FORM}(f)$

TAUTEQ ---: #2#2#2#2#2#2# | [f 1 ← f], 1: -;

```

UNIFY ----:#2#2#2#2#2#2 , -;
T A U T E Q -----:#1 , 1:-;

∀e PROOF f cc sq ;
LABEL GENE 1;
TAUTEQ PROOFTREE(sq)∧INDVAR(x)∧TERM(x)∧(GENI(f cc sq,sq,x,x) v --: v
      EXI(f cc sq,sq,x,x)) 1:-;
UNIFY --:#2#2#2#1 , - ;
LABEL PROOFTR;
TAUT ---:#1 , 1:-;

```

```

∧e HPAUX :#2#2;
∀e - | f1;

∀e DEFEND f cc sq, sq, f 1;
∧E GENE1 :#2;
UNIFY --:#2#2#2#2#2, - ;

```

```

TAUTEQ DEPEND(f cc sq,f1)⊃ AXIOM (f1),1:-;
Vi - ,f1←f1; --
TAUTEQ f=scar(f cc sq) 1:-;
∧i PROOFTR, - , - ;
LABEL USEFUL;
∀e PROVABLE f;
UNIFY -:#2#2,--;
TAUT --:#1 , 1:-;
LABEL C1TH1;
⇒I HPT,-;

```

Proof of the Lemma $\text{BEW}(f) \supset \text{BEW}(x \text{ gen } f)$ Under the Assumption: $\text{INDVAR}(x) \wedge \text{FORM}(f)$

```

LABEL HPT1;
ASSUME BEW(f);

TAUT USEFUL:@2 , -,HPT 1 ,USEFUL;
∧E -:#2
∃e - ,sq;

∧e -:#2#2;
∀e - , f1;
∀e GENRUL2 x,sq;
∀e THEORY x,f1;
TAUTEQ --:#2#1#2#1 [f←f1],HPTT,HPT1:-;
Vi - ,f1←f1;
TAUT -----:#1 ,HPTT,HPT1:-;

∀e GENRUL 1 ((x gen f) cc sq) , sq ,x,x ;
LABEL THN2;
∀e THEON1 x gen f , sq ;
∀e THEON2 x gen f , sq ;
V E TH1 x ,f;
VE TH2 x gen f;
VE TH5 sq;
TAUTEQ -----:#2#2#2#1 [f1 ← f] ,HPTT, THTAUT,HPT1:-;

```

```

UNIFY -----: #2#2#2 , -;
VE TH3, x gen f, sq;
TAUTEQ -----: #1, HPTT, THTAUT, HPT1:-; .

```

```

Ve PROOF (x gen f) cc sq ;
Ve TH4, x;
LABEL GEN1;
TAUTEQ PROOFTREE(sq) ^ INDVAR(x) ^ TERM(x) ^ (---: v GENE((x gen f) cc sq, sq, x, x) v
EXI((x gen f) cc sq, sq, x, x)) HPTT, HPT1:-;
UNIFY ---: #2#2#2#1 , -;

```

```

LABEL PROOFTRI ;
TAUT ---: #1, HPT1:-, THTAUT, HPTT;

```

```

Ve DEPEND (x gen f) cc sq, sq, f 1;
^E GEN1: #2;
3i - , x ← f OCC 2 5 8 11;
3i - , x ← x1 OCC 1 3 5 7;

```

```

TAUTEQ DEPEND((x gen f) cc sq, f 1) ⊃ AXIOM (f 1) , THTAUT, HPTT, HPT1:-;
Vi -, f1 ← f1;
TAUTEQ x gen f = scar((x gen f) cc sq), HPTT, HPT1:-;
A1 PROOFTRI, -, -;
Ve PROVABLE x gen f;
UNIFY -: #2#2, -;
TAUT -: #1, THTAUT, HPT1:-;
LABEL C2TH1;
⊃1 HPT1, -;
≡1 C1 TH1, C2TH1;
LABEL THGEN;
⊃1 HPTT, -;
V1 -, x, f;
Ve TH1 x1, x2 gen f;
Ve TH1 x2, f;
Ve TH1 x1, f;
- Ve TH1 x2, x1 gen f;
VE TH1, x1, f;
VE TH1, x2, f;
TAUT (INDVAR(x1) ^ (INDVAR(x2) ^ FORM(f))) ⊃ (BEW(x1 gen (x2 gen f))) •
BEW(x2 gen (x1 gen f)), THGEN:-;
V1 -, x1, x2, f;

```

5.6 Printout of the proof in the earlier axiomatization

- 1 $\forall x f. ((\text{INDVAR}(x) \wedge \text{FORM}(f)) \supset \text{sbt}(x, x, f) = f)$ (1) ASSUME
- 2 $\forall s sq. ((\text{STRING}(s) \wedge \text{SEQUENCE}(sq)) \supset \text{scar}(s \text{ cc } sq) = s)$ (2) ASSUME
- 3 $\forall s sq. ((\text{STRING}(s) \wedge \text{SEQUENCE}(sq)) \supset \text{s cdr}(s \text{ cc } sq) = sq)$ (3) ASSUME

```

4  $\forall x. ((\text{INDVAR}(x) \wedge \text{FORM}(f)) \Rightarrow \text{FORM}(x_{\text{gen}} f))$  (4) ASSUME
5  $\forall f. (\text{FORM}(f) \Rightarrow \text{STRING}(f))$  (5) ASSUME
6  $\forall f \text{ sq}. ((\text{FORM}(f) \wedge \text{SEQUENCE}(\text{sq})) \Rightarrow \text{SEQUENCE}(f_{\text{cc}} \text{ sq}))$  (6) ASSUME
7  $\forall x. (\text{INDVAR}(x) \Rightarrow \text{TERM}(x))$  (7) ASSUME
8  $\forall pf. (\text{PROOFTREE}(pf) \Rightarrow \text{SEQUENCE}(pf))$  (8) ASSUME
9  $\text{INDVAR}(x) \wedge \text{FORM}(f)$  (9) ASSUME
10 BEW( $x_{\text{gen}} f$ ) (10) ASSUME
11  $(\text{INDVAR}(x) \wedge \text{FORM}(f)) \Rightarrow \text{sbt}(x, x, f) = f$  (1)  $\forall e 1 x, f$ 
12 BEW( $x_{\text{gen}} f$ )  $\equiv (\text{FORM}(x_{\text{gen}} f) \wedge \exists \text{sq}. (\text{PROOFTREE}(\text{sq}) \wedge ((x_{\text{gen}} f) = \text{scar}(\text{sq}) \wedge \forall f1. (\text{DEPEND}(\text{sq}, f1) \Rightarrow \text{AXIOM}(f1))))))$   $\forall e$  PROVABLE  $x_{\text{gen}} f$ 
13  $(\text{INDVAR}(x) \wedge \text{FORM}(f)) \Rightarrow \text{FORM}(x_{\text{gen}} f)$  (4)  $\forall e 4 x, f$ 
14  $\exists \text{sq}. (\text{PROOFTREE}(\text{sq}) \wedge ((x_{\text{gen}} f) = \text{scar}(\text{sq}) \wedge \forall f1. (\text{DEPEND}(\text{sq}, f1) \Rightarrow \text{AXIOM}(f1))))$  (1 4 9 10) g : 13
15  $\text{FORM}(f) \Rightarrow \text{STRING}(f)$  (5)  $\forall e 5 f$ 
16  $(\text{FORM}(f) \wedge \text{SEQUENCE}(\text{sq})) \Rightarrow \text{SEQUENCE}(f_{\text{cc}} \text{ sq})$  (6)  $\forall e 6 f, \text{sq}$ 
17  $\text{INDVAR}(x) \Rightarrow \text{TERM}(x)$  (7)  $\forall e 7 x$ 
18  $\text{PROOFTREE}(\text{sq}) \Rightarrow \text{SEQUENCE}(\text{sq})$  (8)  $\forall e 8 \text{sq}$ 
19  $\text{PROOFTREE}(\text{sq}) \wedge ((x_{\text{gen}} f) = \text{scar}(\text{sq}) \wedge \forall f1. (\text{DEPEND}(\text{sq}, f1) \Rightarrow \text{AXIOM}(f1)))$  (19) ASSUME
20 GENE( $f_{\text{cc}} \text{sq}, \text{sq}, x, x$ )  $\equiv (\text{SEQUENCE}(f_{\text{cc}} \text{sq}) \wedge (\text{INDVAR}(x) \wedge (\text{TERM}(x) \wedge (\text{schr}(f_{\text{cc}} \text{sq}) = \text{sq} \wedge (\text{PROOFTREE}(\text{sq}) \wedge \exists f1. (\text{FORM}(f1) \wedge (\text{scar}(\text{sq}) = (x_{\text{gen}} f1) \wedge \text{scar}(f_{\text{cc}} \text{sq}) = \text{sbt}(x, x, f1))))))))$   $\forall e$  GENRULO  $f_{\text{cc}} \text{sq}, \text{sq}, x, x$ 
21  $(\text{STRING}(f) \wedge \text{SEQUENCE}(\text{sq})) \Rightarrow \text{scar}(f_{\text{cc}} \text{sq}) = f$  (2)  $\forall e 2 f, \text{sq}$ 
22  $(\text{STRING}(f) \wedge \text{SEQUENCE}(\text{sq})) \Rightarrow \text{schr}(f_{\text{cc}} \text{sq}) = \text{sq}$  (3)  $\forall e 3 f, \text{sq}$ 
23  $\text{FORM}(f) \wedge (\text{scar}(\text{sq}) = (x_{\text{gen}} f) \wedge \text{scar}(f_{\text{cc}} \text{sq}) = \text{sbt}(x, x, f))$  (1 2 3 4 5 6 7 8 9 10 19) 1 : 22
24  $\exists f1. (\text{FORM}(f1) \wedge (\text{scar}(\text{sq}) = (x_{\text{gen}} f1) \wedge \text{scar}(f_{\text{cc}} \text{sq}) = \text{sbt}(x, x, f1)))$  (1 2 3 4 5 6 7 8 9 10 19) UNIFY 23
25 GENE( $f_{\text{cc}} \text{sq}, \text{sq}, x, x$ ) (1 2 3 4 5 6 7 8 9 10 19) 1 : 24
26  $\text{PROOFTREE}(f_{\text{cc}} \text{sq}) \equiv ((\text{SEQUENCE}(f_{\text{cc}} \text{sq}) \wedge \text{FORM}(f_{\text{cc}} \text{sq})) \vee (\exists pf. (\text{PROOFTREE}(pf) \wedge (\text{ORI}(f_{\text{cc}} \text{sq}, pf) \vee (\text{ANDE}(f_{\text{cc}} \text{sq}, pf) \vee (\text{FALSEE}(f_{\text{cc}} \text{sq}, pf) \vee (\text{NOTI}(f_{\text{cc}} \text{sq}, pf) \vee (\text{NOTE}(f_{\text{cc}} \text{sq}, pf) \vee (\text{IMPLI}(f_{\text{cc}} \text{sq}, pf)))))))))) \vee (\exists pf \ x \ t. (\text{PROOFTREE}(pf) \wedge (\text{INDVAR}(x) \wedge (\text{TERM}(t) \wedge (\text{GENI}(f_{\text{cc}} \text{sq}, pf, x, t) \vee (\text{GENE}(f_{\text{cc}} \text{sq}, pf, x, t) \vee (\text{EXI}(f_{\text{cc}} \text{sq}, pf, x, t)))))) \vee (\exists pf1 \ pf2. (\text{PROOFTREE}(pf1) \wedge (\text{PROOFTREE}(pf2) \wedge (\text{ANDI}(f_{\text{cc}} \text{sq}, pf1, pf2) \vee$ 

```

(FALSE!(f cc sq, pf1, pf2) v IMPL (f cc sq, pf1, pf2)))) v (3 pf1 pf2 x1 x2. (PROOFTREE(pf1) ^
 (PROOFTREE(pf2) ^ (INDVAR(x1) ^ (INDVAR(x2) ^ EXE(f cc sq, pf1, pf2, x1, x2)))))) v 3 pf1 pf2 pf3.
 (PROOFTREE(pf1) ^ (PROOFTREE(pf2) ^ (PROOFTREE(pf3) ^ ORE(f cc sq, pf1, pf2, pf3))))))
 VE PROOF f cc sq

27 PROOFTREE(sq) ^ (INDVAR(x) ^ (TERM(x) ^ (GENI(f cc sq, sq, x, x) v (GENE(f cc sq, sq, x, x) v
 EXI(f cc sq, sq, x, x)))))) (1 2 3 4 5 6 7 8 9 10 19) 1 : 26

28 3 pf x t. (PROOFTREE(pf) ^ (INDVAR(x) ^ (TERM(t) ^ (GENI(f cc sq, pf, x, t) v (GENE(f cc sq, pf, x, t) v
 EXI(f cc sq, pf, x, t)))))) (1 2 3 4 5 6 7 8 9 10 19) UNIFY 27

29 PROOFTREE(f cc sq) (1 2 3 4 5 6 7 8 9 10 19) 1 : 28

30 V f 1. (DEPEND(sq, f 1) => AXIOM(f 1)) (19) ^E 19 : #2#2

31 DEPEND(sq, f 1) => AXIOM(f 1) (19) VE 30 f 1

32 ((PROOFTREE(f cc sq) ^ (PROOFTREE(sq) ^ sq = scdr(f cc sq))) => (DEPEND(f cc sq, f 1) = DEPEND(sq, f 1))) =
 (ORI(f cc sq, sq) v (ANDE(f cc sq, sq) v (FALSE!(f cc sq, sq) v (3 f. (FORM(f) ^ ((NOTID(f cc sq, sq, f) v
 (NOTED(f cc sq, sq, f) v IMPLID(f cc sq, sq, f))) ^ f, f 1)) v 3 x t. (INDVAR(x) ^ (TERM(t) ^
 (GENI(f cc sq, sq, x, t) v (GENE(f cc sq, sq, x, t) v EXI(f cc sq, sq, x, t))))))))))
 VE DEPEND f cc sq, sq, f 1

33 INDVAR(x) ^ (TERM(x) ^ (GENI(f cc sq, sq, x, x) v (GENE(f cc sq, sq, x, x) v EXI(f cc sq, sq, x, x))))
 (I 23 4 5 6 7 8 9 10 19) ^E 27 : #2

34 3 x t. (INDVAR(x) ^ (TERM(t) ^ (GENI(f cc sq, sq, x, t) v (GENE(f cc sq, sq, x, t) v EXI(f cc sq, sq, x, t))))))
 (I 2 3 4 5 6 7 8 9 10 19) UNIFY 33

35 DEPEND(f cc sq, f 1) => AXIOM(f 1) (1 2 3 4 5 6 7 8 9 10 19) 1 : 34

36 V f 1. (DEPEND(f cc sq, f 1) => AXIOM(f 1)) (1 2 3 4 5 6 7 8 9 10 19) VI 35 f 1 + f 1

37 f = scar(f cc sq) (1 2 3 4 5 6 7 8 9 10 19) 1 : 36

38 PROOFTREE(f cc sq) ^ (f = scar(f cc sq) ^ V f 1. (DEPEND(f cc sq, f 1) => AXIOM(f 1)))
 (1 2 3 4 5 6 7 8 9 10 19) AI (29 (37 36))

39 BEW(f) = (FORM(f) ^ 3 sq. (PROOFTREE(sq) ^ (f = scar(sq) ^ V f 1. (DEPEND(sq, f 1) => AXIOM(f 1))))
 VE PROVABLE f

40 3 sq. (PROOFTREE(sq) ^ (f = scar(sq) ^ V f 1. (DEPEND(sq, f 1) => AXIOM(f 1)))) (1 2 3 4 5 6 7 8 9 10) UNIFY 38

41 BEW(f) (1 2 3 4 5 6 7 8 9 10) 9, 39, 40

42 BEW(x gen f) => BEW(f) (1 2 3 4 5 6 7 8 9) => 10 41

43 BEW(f) (43) ASSUME

44 FORM(f) ^ 3 sq. (PROOFTREE(sq) ^ (f = scar(sq) ^ V f 1. (DEPEND(sq, f 1) => AXIOM(f 1)))) (43) 43, 43, 39

45 3 sq. (PROOFTREE(sq) ^ (f = scar(sq) ^ V f 1. (DEPEND(sq, f 1) => AXIOM(f 1)))) (43) ^E 44 : #2

```

46 PROOFTREE(sq)^(f=scar(sq)^(Vf1.(DEPEND(sq,f1)=AXIOM(f1))) (46) ASSUME
47 Vf1.(DEPEND(sq,f1)=AXIOM(f1)) (46) ^E 46 :#2*2
48 DEPEND(sq,f1)=AXIOM(f1) (46) VE 47 f1
49 APGENI(x,sq)=((INDVAR(x)^(Vf1.(DEPEND(sq,f)=~FR(x,f)))^PROOFTREE(sq)) VE GENRUL2 x , sq
50 AXIOM(f1)=~FR(x,f1)^FORM(f1) VE THEORY x , f1
51 DEPEND(sq,f1)=~FR(x,f1) (1 2 3 4 5 6 7 8 9 43 46) 9 , 43 : 50
52 Vf1.(DEPEND(sq,f1)=~FR(x,f1)) (1 2 3 4 5 6 7 8 9 43 46) VI 51 f1 ← f1
53 APGENI(x,sq) (1 2 3 4 5 6 7 8 9 43 46) 9 , 43 : 52
54 GENI((x gen f) cc sq,sq,x,x) (SEQUENCE((x gen f) cc sq)^(INDVAR(x)^(INDVAR(x)^(
  scdr((x gen f) cc sq)=sq^(PROOFTREE(sq)^(Vf1.(FORM(f1)^(scar((x gen f) cc sq)=
  (x gen f1)^(scar(sq)=sbt(x,x,f1)^APGENI(x,sq))))))))))
  VE GENRUL1(x gen f) cc sq , sq , x , x
55 (STRING(x gen f)^SEQUENCE(sq))=scar((x gen f) cc sq)=(x gen f)(2) VE 2 x gen f , sq
56 (STRING(x gen f)^SEQUENCE(sq))=scdr((x gen f) cc sq)=sq (3) VE 3 x gen f , sq
57 (INDVAR(x)^FORM(f))=FORM(x gen f) (4) VE 4 x , f
58 FORM(x gen f)=STRING(x gen f) (5) VE 5 x gen f
59 PROOFTREE(sq)=SEQUENCE(sq) (8) VE 8 sq
60 FORM(f)^(scar((x gen f) cc sq)=(x gen f)^(scar(sq)=sbt(x,x,f)^APGENI(x,sq)))
  (1 2 3 4 5 6 7 8 9 43 46) 11 , 43 : 59 , 9
61 V f1.(FORM(f1)^(scar((x gen f) cc sq)=(x gen f1)^(scar(sq)=sbt(x,x,f1)^
  APGENI(x,sq)))) (1 2 3 4 5 6 7 8 9 43 46) UNIFY 60
62 (FORM(x gen f)^SEQUENCE(sq))=SEQUENCE((x gen f) cc sq) (6) VE 6 x gen f , sq
63 GENI((x gen f) cc sq,sq,x,x) (1 2 3 4 5 6 7 8 9 43 46) 9 , 11 , 43 : 62
64 PROOFTREE((x gen f) cc sq)^(SEQUENCE((x gen f) cc sq)^FORM((x gen f) cc sq))^(
  (Vpf.(PROOFTREE(pf)^(ORI((x gen f) cc sq,pf)^(ANDE((x gen f) cc sq,pf)^(
  FALSEE((x gen f) cc sq,pf)^(NOTI((x gen f) cc sq,pf)^(NOTE((x gen f) cc sq,pf)^(
  IMPLI((x gen f) cc sq,pf))))))^(Vpf x1t.(PROOFTREE(pf)^(INDVAR(x1)^(TERM(t)^(
  GENI((x gen f) cc sq,pf,x1,t)^(GENE((x gen f) cc sq,pf,x1,t)^(EXI((x gen f)
  cc sq,pf,x1,t))))))^(Vpf1pf2.(PROOFTREE(pf1)^(PROOFTREE(pf2)^(ANDI((x gen f)
  cc sq,pf1,pf2)^(FALSEI((x gen f) cc sq,pf1,pf2)^(IMPLE((x gen f) cc sq,pf1,pf2))))))^(
  (3pf1pf2x1x2.(PROOFTREE(pf1)^(PROOFTREE(pf2)^(INDVAR(x1)^(INDVAR(x2)^(EXE(
  (x gen f) cc sq,pf1,pf2,x1,x2))))))^(V3pf1pf2pf3.(PROOFTREE(pf1)^(PROOFTREE(pf2)^(
  PROOFTREE(pf3)^(ORE((x gen f) cc sq,pf1,pf2,pf3)))))))))) VE PROOF (x gen f) cc sq
65 INDVAR(x)=TERM(x) (7) VE 7 x

```

6 6 **PROOFTREE**(sq) $\wedge$ (**INDVAR**(x) $\wedge$ (**TERM**(x) $\wedge$ (**GENI**((x gen f) cc sq,sq,x,x) $\vee$ (**GENE**((x gen f) cc sq,sq,x,x) $\vee$ **EXI**((x gen f) cc sq,sq,x,x)))) (1 2 3 4 5 6 7 8 9 43 46) 9,43 : 65
 6 7 $\exists$ pf x1 t.(**PROOFTREE**(pf) $\wedge$ (**INDVAR**(x1) $\wedge$ (**TERM**(t) $\wedge$ (**GENI**((x gen f) cc sq,pf,x1,t) $\vee$ (**GENE**((x gen f) cc sq,pf,x1,t) $\vee$ **EXI**((x gen f) cc sq,pf,x1,t)))))) (1 2 3 4 5 6 7 8 9 43 46) UNIFY 66
 68 **PROOFTREE**((x gen f) cc sq) (1 2 3 4 5 6 7 8 9 43 46) 43 : 67 , 11 , 9
 69 ((**PROOFTREE**((x gen f) cc sq) $\wedge$ (**PROOFTREE**(sq) $\wedge$ sq=scdr((x gen f) cc sq))) $\Rightarrow$ (**DEPEND**((x gen f) cc sq,f1) $\Rightarrow$ **DEPEND**(sq,f1)) $\Rightarrow$ (**ORI**((x gen f) cc sq,sq) $\vee$ (**ANDE**((x gen f) cc sq,sq) $\vee$ (**FALSEE**((x gen f) cc sq,sq) $\vee$ ($\exists$ f.(**FORM**(f) $\wedge$ (**NOTID**((x gen f) cc sq,sq,f) $\vee$ (**NOTED**((x gen f) cc sq,sq,f) $\vee$ **IMPLID**((x gen f) cc sq,sq,f))) $\wedge$ f $\neq$ f1)) $\vee$ $\exists$ x1 t.(**INDVAR**(x1) $\wedge$ (**TERM**(t) $\wedge$ (**GENI**((x gen f) cc sq,sq,x1,t) $\vee$ (**GENE**((x gen f) cc sq,sq,x1,t) $\vee$ **EXI**((x gen f) cc sq,sq,x1,t)))))) $\vee$ **DEPEND** (x gen f) cc sq , sq , f1
 7 0 **INDVAR**(x) $\wedge$ (**TERM**(x) $\wedge$ (**GENI**((x gen f) cc sq,sq,x,x) $\vee$ (**GENE**((x gen f) cc sq,sq,x,x) $\vee$ **EXI**((x gen f) cc sq,sq,x,x)))) (1 2 3 4 5 6 7 8 9 43 46) $\wedge$ E 66 :#2
 7 1 $\exists$ t.(**INDVAR**(x) $\wedge$ (**TERM**(t) $\wedge$ (**GENI**((x gen f) cc sq,sq,x,t) $\vee$ (**GENE**((x gen f) cc sq,sq,x,t) $\vee$ **EXI**((x gen f) cc sq,sq,x,t)))) (1 2 3 4 5 6 7 8 9 43 46) 70 x $\leftarrow$ t OCC
 7 2 $\exists$ x1 t.(**INDVAR**(x1) $\wedge$ (**TERM**(t) $\wedge$ (**GENI**((x gen f) cc sq,sq,x1,t) $\vee$ (**GENE**((x gen f) cc sq,sq,x1,t) $\vee$ **EXI**((x gen f) cc sq,sq,x1,t)))) (1 2 3 4 5 6 7 8 9 43 46) 71 x $\leftarrow$ x1 OCC
 73 **DEPEND**((x gen f) cc sq,f1) $\Rightarrow$ **AXIOM**(f1) (1 2 3 4 5 6 7 8 9 43 46) 11 , 9 , 43 : 72
 74 $\forall$ f1.(**DEPEND**((x gen f) cc sq,f1) $\Rightarrow$ **AXIOM**(f1)) (1 2 3 4 5 6 7 8 9 43 46) VI 73 f1 $\leftarrow$ f1
 75 (x gen f)=scar((x gen f) cc sq) (1 2 3 4 5 6 7 8 9 43 46) 9 , 43 : 74
 7 6 **PROOFTREE**((x gen f) cc sq) $\wedge$ ((x gen f)=scar((x gen f) cc sq) $\wedge$ $\forall$ f1.(**DEPEND**((x gen f) cc sq,f1) $\Rightarrow$ **AXIOM**(f1))) (1 2 3 4 5 6 7 8 9 43 46) $\wedge$ I (68 (75 74))
 77 **BEW**(x gen f) $\Rightarrow$ (**FORM**(x gen f) $\wedge$ $\exists$ sq.(**PROOFTREE**(sq) $\wedge$ ((x gen f)=scar(sq) $\wedge$ $\forall$ f1.(**DEPEND**(sq,f1) $\Rightarrow$ **AXIOM**(f1)))) $\vee$ **PROVABLE** x gen f
 7 8 $\exists$ sq.(**PROOFTREE**(sq) $\wedge$ ((x gen f)=scar(sq) $\wedge$ $\forall$ f1.(**DEPEND**(sq,f1) $\Rightarrow$ **AXIOM**(f1)))) (1 2 3 4 5 6 7 8 9 10 19 43 46) UNIFY 76
 79 **BEW**(x gen f) (1 2 3 4 5 6 7 8 9 43) 11 , 9 , 43 : 78
 8 0 **BEW**(f) $\Rightarrow$ **BEW**(x gen f) (1 2 3 4 5 6 7 8 9) $\Rightarrow$ I 43 79
 81 **BEW**(x gen f) $\Rightarrow$ **BEW**(f) (1 2 3 4 5 6 7 8 9) $\Rightarrow$ I 42 80
 8 2 (**INDVAR**(x) $\wedge$ **FORM**(f)) $\Rightarrow$ (**BEW**(x gen f) $\Rightarrow$ **BEW**(f)) (1 2 3 4 5 6 7 8) $\Rightarrow$ I 9 81
 8 3 $\forall$ x f.((**INDVAR**(x) $\wedge$ **FORM**(f)) $\Rightarrow$ (**BEW**(x gen f) $\Rightarrow$ **BEW**(f))) (1 2 3 4 5 6 7 8) VI 82 x , f
 8 4 (**INDVAR**(x1) $\wedge$ **FORM**(x2 gen f)) $\Rightarrow$ (**BEW**(x1 gen (x2 gen f)) $\Rightarrow$ **BEW**(x2 gen f)) (1 2 3 4 5 6 7 8) $\vee$ E 83 x1 , x2 gen f

85 $(\text{INDVAR}(x_2) \wedge \text{FORM}(f)) \supset (\text{BEW}(x_2 \text{ gen } f) \equiv \text{BEW}(f))$ (1 2 3 4 5 6 7 8) VE 83 x_2, f

86 $(\text{INDVAR}(x_1) \wedge \text{FORM}(f)) \supset (\text{BEW}(x_1 \text{ gen } f) \equiv \text{BEW}(f))$ (1 2 3 4 5 6 7 8) VE 83 x_1, f

87 $(\text{INDVAR}(x_2) \wedge \text{FORM}(x_1 \text{ gen } f)) \supset (\text{BEW}(x_2 \text{ gen } (x_1 \text{ gen } f)) \equiv \text{BEW}(x_1 \text{ gen } f))$
(1 2 3 4 5 6 7 8) VE 83 $x_2, x_1 \text{ gen } f$

88 $(\text{INDVAR}(x_1) \wedge \text{FORM}(f)) \supset \text{FORM}(x_1 \text{ gen } f)$ (4) VE 4 x_1, f

89 $(\text{INDVAR}(x_2) \wedge \text{FORM}(f)) \supset \text{FORM}(x_2 \text{ gen } f)$ (4) VE 4 x_2, f

90 $(\text{INDVAR}(x_1) \wedge (\text{INDVAR}(x_2) \wedge \text{FORM}(f))) \supset (\text{BEW}(x_1 \text{ gen } (x_2 \text{ gen } f)) \equiv \text{BEW}(x_2 \text{ gen } (x_1 \text{ gen } f)))$
(1 2 3 4 5 6 7 8) 84 : 89

91 $\forall x_1 x_2 f. ((\text{INDVAR}(x_1) \wedge (\text{INDVAR}(x_2) \wedge \text{FORM}(f))) \supset (\text{BEW}(x_1 \text{ gen } (x_2 \text{ gen } f)) \equiv \text{BEW}(x_2 \text{ gen } (x_1 \text{ gen } f))))$ (1 2 3 4 5 6 7 8) VI 90 x_1, x_2, f

REFERENCES

Prawitz, D.,

1965 *Natural deduction, A proof theoretical study*
Almqvist and Wiksell, Stockholm (1965).

Godel, K.,

1930 *Die Vollständigkeit der Axiome des logischen Funktionenkalküls*
Monatshefte für Mathematik und Physik 37, (1930) 349-360.

Godel, K.,

1931 *Über formal unentscheidbare Sätze der Principia mathematica und verwandter Systeme I*, Monatshefte für Mathematik und Physik 38, (1931) 173-198.

Weyhrauch, R.W., and Thomas, A.J.,

1974 **FOL: A Proof Checker for First-order Logic**
Stanford Artificial Intelligence Laboratory, Memo AIM-235 (1974).

